

ULAM STABILITY FOR FIRST-ORDER NONLINEAR DYNAMIC EQUATIONS

MARTIN BOHNER AND SANKET TIKARE

Dedicated to Professor Mustafa Kulenović on the occasion of his 70th birthday

ABSTRACT. The purpose of this paper is to investigate Ulam stability of first-order nonlinear dynamic equations on time scales. Based on the method of the Picard operator and using dynamic inequalities, we obtain four types of stability. In addition, as applications of our main result, we obtain new Ulam stability results for other nonlinear dynamic equations. An example is also provided to illustrate our main result.

1. INTRODUCTION

It is widely known that stability of solutions is one of the most important and interesting properties among various qualitative properties of solutions. In the existing literature, there are several stability theories, for both differential and difference equations (see e.g., [15, 20, 23, 24] for the discrete case and [25, 26] for PDEs) but the concept of Ulam stability has significant applications in various fields of mathematical analysis. This is mainly because Ulam stability essentially deals with the existence of an exact solution near to every approximate solution and is useful in the situation when it is difficult to find the exact solution. This kind of stability for functional equations was first discussed by Ulam [39] in his famous talk at the University of Wisconsin in 1940. He proposed to “provide an approximate solution for the exact solution in a simple form for a functional equation”. One year later, D. H. Hyers [22] delivered an affirmative answer to this question. Thereafter, the results of Hyers were extended by many authors, but remarkable improvements were provided separately by T. Aoki [10], D. G. Bourgin [19], and Th. M. Rassias [31]. The problem of stability in Ulam sense for various kinds of differential, difference, integral equations etc. has been seriously studied by many researchers employing several techniques. In 2005, D. Popa [30] studied Ulam-type stability for difference equations. Some very recent studies on Ulam stability for difference equations can be found in [5–8, 11–13, 27]. I. A. Rus [33] presented four types of Ulam stability for differential equations

2010 *Mathematics Subject Classification.* 34N05, 34D20, 39A30, 47H09.

Key words and phrases. Dynamic equations; Ulam stability; Picard operator; Gronwall inequality.

**Certified as
TRUE COPY**



Principal
Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.

$$x'(t) = f(t, x(t))$$

both in finite and infinite intervals. Also, in [34], he presented and discussed Ulam-type stability for the differential equation

$$x'(t) = p(t) + f(t, x(t))$$

in Banach spaces. Y. Shen and Y. Li [36], employing the method of variation of parameters, established the Ulam stability for linear differential equations of first-order, second-order, third-order, and n th order. In 2015, J. Huang et al. [21], adopting a fixed point method, investigated Hyers–Ulam as well as generalized Hyers–Ulam stability of nonlinear differential equations involving a Lipschitz condition on infinite intervals. Q. H. Alqifiary and S. M. Jung [3] proved Hyers–Ulam stability for second-order differential equations using Gronwall's inequality. Very recently, A. B. Makhoulf et al. [14] investigated Hyers–Ulam and Hyers–Ulam–Rassias stability for stochastic functional differential equations via the method of fixed point and stochastic analysis techniques.

S. András and A. R. Mészáros [9] studied Hyers–Ulam stability of some linear and nonlinear dynamic equations as well as integral equations on time scales. They employed both direct and operational methods, and based on the theory of Picard's operators, proposed a unified approach to Hyers–Ulam stability. Y. Shen [35], employing the method of integrating factor, investigated Ulam stability of the first-order linear dynamic equation

$$x^\Delta(t) = p(t)x(t) + f(t)$$

and its adjoint equation

$$x^\Delta(t) = -p(t)x^\sigma(t) + f(t)$$

on a finite interval. Also, D. R. Anderson and M. Ointsuka [4] established Hyers–Ulam stability of certain first-order linear homogeneous dynamic equations with constant coefficients. They extended the results given in [28, 29] to all time scales and also provided an application to a perturbed linear dynamic equation.

Most recently, in 2021, applying dynamic inequalities, M. A. Alghamdi et al. [1, 2] obtained several results on Hyers–Ulam and Hyers–Ulam–Rassias stability for the first-order dynamic equations

$$x^\Delta(t) = p(t)x(t) + f(t)$$

and

$$x^\Delta(t) = p(t)x(t) + f(t, x(t), h(x(t))) + g(t),$$

respectively.

In this paper, we investigate Ulam stability for the nonlinear dynamic equation (NDE) of the form

$$x^\Delta(t) + p(t)x^\sigma(t) = f(t, x(t)), \quad t \in \mathbb{J}^k, \quad (1.1)$$

where $\mathbb{J} := [a, b]_{\mathbb{T}}$, $a, b \in \mathbb{T}$ with $a < b$, $x : \mathbb{J} \rightarrow \mathbb{R}$ is the unknown function to be determined, $x^\sigma = x \circ \sigma$, x^Δ is the delta derivative of x , $p : \mathbb{T} \rightarrow \mathbb{R}$ is a positively

Certified as
TRUE COPY

Principal

Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.

regressive and rd-continuous function, $f : \mathbb{J} \times \mathbb{R} \rightarrow \mathbb{R}$ is rd-continuous in its first variable and continuous in its second variable.

Based on the method of Picard operator and dynamic inequalities, we obtain results on stability of NDE (1.1). The results obtained in this paper are more general than the known results available in the literature and include the studies [2, 4, 21, 33, 36]. For the existence, uniqueness, and other properties of solutions of NDE (1.1), we refer to [18, 37, 38].

2. PRELIMINARIES

To understand the notation used in this paper, we include some preliminary material. The following material pertinent to time scales can be found in [16, 17]. A nonempty closed subset of the real line \mathbb{R} is called a time scale \mathbb{T} . We usually write $\mathbb{T}^\kappa = \mathbb{T} \setminus \{\max \mathbb{T}\}$ if $\max \mathbb{T} < \infty$, otherwise $\mathbb{T}^\kappa = \mathbb{T}$.

Definition 2.1. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is said to be delta differentiable at $t \in \mathbb{T}^\kappa$ if there exists $f^\Delta(t) \in \mathbb{R}$, a so-called delta derivative of f , with the following property: For any $\varepsilon > 0$ there is a neighbourhood N of t , such that

$$|f(\sigma(t)) - f(s) - f^\Delta(t)(\sigma(t) - s)| \leq \varepsilon |\sigma(t) - s| \quad \text{for all } s \in N.$$

Definition 2.2. A function $f : \mathbb{T} \rightarrow \mathbb{R}$ is rd-continuous if it is continuous at every right-dense point or maximal point in \mathbb{T} and its left sided limits exist at left-dense points in \mathbb{T} . The symbol $C_{rd}(\mathbb{T}, \mathbb{R})$ will be used for the set of all such functions. If a function $f : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$ is rd-continuous in its first variable and continuous in its second variable, then we write $f \in C_{rd}(\mathbb{T} \times \mathbb{R}, \mathbb{R})$.

Remark 2.1. The family $C_{rd}(\mathbb{J}, \mathbb{R})$ of all rd-continuous functions from \mathbb{J} into \mathbb{R} forms a Banach space coupled with the norm $\|\cdot\|$ defined as $\|x\| := \sup_{t \in \mathbb{J}} |x(t)|$.

Definition 2.3. We say that $p : \mathbb{T} \rightarrow \mathbb{R}$ is regressive if $1 + \mu(t)p(t) \neq 0$ for all $t \in \mathbb{T}$. The symbol $\mathcal{R}(\mathbb{T}, \mathbb{R})$ will be used for the set of all rd-continuous regressive functions. If $1 + \mu(t)p(t) > 0$ for all $t \in \mathbb{T}$, then p is said to be positively regressive, and $\mathcal{R}^+(\mathbb{T}, \mathbb{R})$ denotes the set of all rd-continuous positively regressive functions.

Definition 2.4. For $p \in \mathcal{R}(\mathbb{T}, \mathbb{R})$, the generalized exponential function $e_p(t, s)$ on the time scale \mathbb{T} is defined as

$$e_p(t, s) := \begin{cases} \exp \left(\int_s^t \frac{\text{Log} |1 + \mu(r)p(r)|}{\mu(r)} \Delta r \right) & \text{if } \mu(r) \neq 0, \\ \exp \left(\int_s^t p(r) \Delta r \right) & \text{if } \mu(r) = 0. \end{cases}$$

For $p, q \in \mathcal{R}(\mathbb{T}, \mathbb{R})$, we define

$$p \oplus q := p + q + \mu p q, \quad \ominus p := \frac{-p}{1 + \mu p}, \quad p \ominus q := p \oplus (\ominus q).$$

**Certified as
TRUE COPY**


Principal

**Ramniranjan Jhunhunwala College,
Ghatkopar (W), Mumbai-400086.**

Remark 2.2. We let

$$E_p := \sup_{s,t \in \mathbb{J}} |e_{\ominus p}(t,s)| > 0 \quad \text{and} \quad E_q := \sup_{s,t \in \mathbb{J}} |e_{\ominus \frac{q}{1-\mu q}}(t,s)| > 0.$$

In our investigation, we mainly use the following results and definition.

Theorem 2.1 (See [40, Theorem 2]). *Let $y, \mathcal{F} \in C_{\text{rd}}(\mathbb{T}, \mathbb{R}^+)$ with \mathcal{F} a nondecreasing function and $\mathcal{G}, \mathcal{H} \in \mathcal{R}^+(\mathbb{T}, \mathbb{R})$ with $\mathcal{G} \geq 0, \mathcal{H} \geq 0$. If*

$$y(t) \leq \mathcal{F}(t) + \int_a^t \mathcal{H}(s) \left[y(s) + \int_a^s \mathcal{G}(\tau) y(\tau) \Delta\tau \right] \Delta s \quad \text{for all } t \in \mathbb{T}^{\kappa},$$

then

$$y(t) \leq \mathcal{F}(t) e_{\mathcal{H} + \mathcal{G}}(t, a) \quad \text{for all } t \in \mathbb{T}^{\kappa}.$$

Definition 2.5 (See [32, Definition 2.1]). *Let (M, d) be a metric space. An operator $A : M \rightarrow M$ is said to be a Picard operator if there exists $u^* \in M$ with the following properties:*

- (i) $F_A = \{u^*\}$, where F_A is the fixed point set of A ;
- (ii) the sequence $\{A^n(u)\}_{n \in \mathbb{N}}$ converges to u^* for all $u \in M$.

Lemma 2.1 (Abstract Gronwall lemma [32, Lemma 2.1]). *Let (M, d, \leq) be an ordered metric space and $A : M \rightarrow M$ an increasing Picard operator ($F_A = u_A^*$). Then for $u \in M$, $u \leq A(s)$ implies $u \leq u_A^*$, while $u \geq A(s)$ implies $u \geq u_A^*$.*

Lemma 2.2 (See [18, Lemma 3.1]). *Let $a \in \mathbb{T}$, $f \in C_{\text{rd}}(\mathbb{J} \times \mathbb{R}, \mathbb{R})$, and $p \in (\mathbb{J}, \mathbb{R})$. Then, x satisfies (1.1) if and only if*

$$x(t) = e_{\ominus p}(t, a)x(a) + \int_a^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \quad \text{for all } t \in \mathbb{J}. \quad (2.1)$$

Now, we introduce some basic definitions that will be used in this paper.

Definition 2.6. *We say that NDE (1.1) has Hyers–Ulam stability if there exists $K > 0$ with the following property: For any $\varepsilon > 0$, if $y \in C_{\text{rd}}^1(\mathbb{J}, \mathbb{R})$ is such that*

$$|y^\Delta(t) + p(t)y^\sigma(t) - f(t, y(t))| \leq \varepsilon \quad \text{for all } t \in \mathbb{J}^{\kappa}, \quad (2.2)$$

then there exists $x \in C_{\text{rd}}^1(\mathbb{J}, \mathbb{R})$ satisfying (1.1) such that

$$|y(t) - x(t)| \leq K\varepsilon \quad \text{for all } t \in \mathbb{J}. \quad (2.3)$$

Such $K > 0$ is known as HUS constant.

Definition 2.7. *We say that NDE (1.1) has generalized Hyers–Ulam stability if there exists $\theta_f \in C(\mathbb{R}^+, \mathbb{R}^+)$, $\theta_f(0) = 0$ with the following property: For any $\varepsilon > 0$, if $y \in C_{\text{rd}}^1(\mathbb{J}, \mathbb{R})$ is such that*

$$|y^\Delta(t) + p(t)y^\sigma(t) - f(t, y(t))| \leq \varepsilon \quad \text{for all } t \in \mathbb{J}^{\kappa}, \quad (2.4)$$

then there exists $x \in C_{\text{rd}}^1(\mathbb{J}, \mathbb{R})$ satisfying (1.1) such that

$$|y(t) - x(t)| \leq \theta_f(\varepsilon) \quad \text{for all } t \in \mathbb{J}. \quad (2.5)$$

Certified as
TRUE COPY

Principal

Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.

Definition 2.8. Let \mathcal{N} be a family of positive nondecreasing rd-continuous functions defined on \mathbb{J} . We say that NDE (1.1) has Hyers–Ulam–Rassias stability of type \mathcal{N} if there exists $K > 0$ with the following property: For any $\phi \in \mathcal{N}$ and $\varepsilon > 0$, if $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ is such that

$$|y^\Delta(t) + p(t)y^\sigma(t) - f(t, y(t))| \leq \varepsilon\phi(t) \quad \text{for all } t \in \mathbb{J}^\kappa, \quad (2.6)$$

then there exists $x \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfying (1.1) such that

$$|y(t) - x(t)| \leq K\varepsilon\phi(t) \quad \text{for all } t \in \mathbb{J}. \quad (2.7)$$

Such $K > 0$ is known as HURS constant.

Definition 2.9. Let \mathcal{N} be a family of positive nondecreasing rd-continuous functions defined on \mathbb{J} . We say that NDE (1.1) has generalized Hyers–Ulam–Rassias stability of type \mathcal{N} if there exists $K > 0$ with the following property: For any $\phi \in \mathcal{N}$, if $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ is such that

$$|y^\Delta(t) + p(t)y^\sigma(t) - f(t, y(t))| \leq \phi(t) \quad \text{for all } t \in \mathbb{J}^\kappa, \quad (2.8)$$

then there exists $x \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfying (1.1) such that

$$|y(t) - x(t)| \leq K\phi(t) \quad \text{for all } t \in \mathbb{J}. \quad (2.9)$$

Such $K > 0$ is known as GHURS constant.

Remark 2.3. A function $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfies (2.6) if there exists $\psi \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ (which depends on y) with the following properties:

- (i) $|\psi(t)| \leq \varepsilon\phi(t)$ for all $t \in \mathbb{J}$,
- (ii) $y^\Delta(t) + p(t)y^\sigma(t) = f(t, y(t)) + \psi(t)$ for all $t \in \mathbb{J}^\kappa$.

Similar arguments hold for the inequalities (2.4) and (2.8).

3. ULAM STABILITY

In this section we prove our main result of Ulam stability for NDE (1.1) and provide its applications.

Theorem 3.1. Consider the NDE (1.1). Assume that the following conditions are satisfied.

(C₁) Let $p \in \mathcal{R}^+(\mathbb{J}, \mathbb{R})$ and $f \in C_{rd}(\mathbb{J} \times \mathbb{R}, \mathbb{R})$.

(C₂) There exists $L_f > 0$ such that

$$|f(t, u) - f(t, v)| \leq L_f|u - v| \quad \text{for all } t \in \mathbb{J} \text{ and } u, v \in \mathbb{R}. \quad (3.1)$$

(C₃) There exists $\eta > 0$ such that for $\phi \in \mathcal{N}(\mathbb{J}, \mathbb{R}^+)$

$$\int_a^t \phi(s)\Delta s \leq \eta\phi(t) \quad \text{for all } t \in \mathbb{J}. \quad (3.2)$$

If $E_p L_f (b - a) < 1$, then the following assertions hold:

**Certified as
TRUE COPY**



Principal

**Ramniranjan Jhunjunwala College,
Ghatkopar (W), Mumbai-400086.**

- (i) The NDE (1.1) has a unique solution $x \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfying the initial condition $x(a) = A$ for any initial value $A \in \mathbb{R}$.
- (ii) The NDE (1.1) has Hyers–Ulam–Rassias stability of type \mathcal{N} with HURS constant $E_p(b-a)e_{E_p L_f}(b, a)$.

Proof. By Lemma 2.2, the NDE (1.1) with initial condition $x(a) = A$ is equivalent to the integral equation

$$x(t) = e_{\ominus p}(t, a)A + \int_a^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \quad \text{for all } t \in \mathbb{J}. \quad (3.3)$$

We first show (i). Fix $A \in \mathbb{R}$ and define $T : C_{rd}(\mathbb{J}, \mathbb{R}) \rightarrow C_{rd}(\mathbb{J}, \mathbb{R})$ by

$$T(x)(t) := e_{\ominus p}(t, a)A + \int_a^t e_{\ominus p}(t, s)f(s, x(s))\Delta s. \quad (3.4)$$

We show that the operator T has a fixed point, and for this we use the contraction mapping principle. For any $x, y \in C_{rd}(\mathbb{J}, \mathbb{R})$, we can write

$$\begin{aligned} & |T(x)(t) - T(y)(t)| \\ & \leq |e_{\ominus p}(t, a)||A - A| + \int_a^t |e_{\ominus p}(t, s)||f(s, x(s)) - f(s, y(s))|\Delta s \\ & \stackrel{(C_2)}{\leq} E_p \int_a^t L_f |x(s) - y(s)|\Delta s \\ & \leq E_p L_f (b-a) \|x - y\|. \end{aligned}$$

Thus,

$$\|T(x) - T(y)\| \leq E_p L_f (b-a) \|x - y\| \quad \text{for all } x, y \in C_{rd}(\mathbb{J}, \mathbb{R}).$$

Since $E_p L_f (b-a) < 1$, the above inequality implies that the operator T is a contraction on $C_{rd}(\mathbb{J}, \mathbb{R})$. So, T has a unique fixed point $x^* \in C_{rd}(\mathbb{J}, \mathbb{R})$, which is the unique solution of the NDE (1.1) satisfying $x^*(a) = A$.

Now we show (ii). Let $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfy (2.6) and let $x \in C_{rd}(\mathbb{J}, \mathbb{R})$ be the unique solution of (1.1) satisfying the initial condition $x(a) = y(a)$. Then (C₁) allows to write

$$x(t) = e_{\ominus p}(t, a)y(a) + \int_a^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \quad \text{for all } t \in \mathbb{J}.$$

Now, since $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfies (2.6), by Remark 2.3, we can write

$$y^\Delta(t) + p(t)y^\sigma(t) = f(t, y(t)) + \psi(t) \quad \text{for all } t \in \mathbb{J}^\kappa,$$

where $\|\psi(t)\| \leq \varepsilon \phi(t)$ for all $t \in \mathbb{J}$. Thus,

$$\begin{aligned} y(t) &= e_{\ominus p}(t, a)y(a) + \int_a^t e_{\ominus p}(t, s)(f(s, y(s)) + \psi(s))\Delta s \\ &= e_{\ominus p}(t, a)y(a) + \int_a^t e_{\ominus p}(t, s)f(s, y(s))\Delta s + \int_a^t e_{\ominus p}(t, s)\psi(s)\Delta s. \end{aligned}$$

This gives

**Certified as
TRUE COPY**

Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

$$\begin{aligned}
 & \left| y(t) - e_{\ominus p}(t, a)y(a) - \int_a^t e_{\ominus p}(t, s)f(s, y(s))\Delta s \right| \\
 & \leq \int_a^t |e_{\ominus p}(t, s)|\psi(s)\Delta s \\
 & \stackrel{(C_3)}{\leq} E_p(b-a)\eta\phi(t)\varepsilon \quad \text{for all } t \in \mathbb{J}.
 \end{aligned} \tag{3.5}$$

Now, for $t \in \mathbb{J}$, we can write

$$\begin{aligned}
 |y(t) - x(t)| &= \left| y(t) - e_{\ominus p}(t, a)y(a) - \int_a^t e_{\ominus p}(t, s)f(s, y(s))\Delta s \right. \\
 & \quad \left. + \int_a^t e_{\ominus p}(t, s)f(s, y(s))\Delta s - \int_a^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \right| \\
 & \leq \left| y(t) - e_{\ominus p}(t, a)y(a) - \int_a^t e_{\ominus p}(t, s)f(s, y(s))\Delta s \right| \\
 & \quad + \int_a^t |e_{\ominus p}(t, s)||f(s, y(s)) - f(s, x(s))|\Delta s \\
 & \stackrel{(3.5)}{\leq} E_p(b-a)\eta\phi(t)\varepsilon + \int_a^t |e_{\ominus p}(t, s)||f(s, y(s)) - f(s, x(s))|\Delta s \\
 & \stackrel{(C_2)}{\leq} E_p(b-a)\eta\phi(t)\varepsilon + L_f \int_a^t |e_{\ominus p}(t, s)(y(s) - x(s))|\Delta s.
 \end{aligned} \tag{3.6}$$

According to (3.6), we consider the operator $S : C_{rd}(\mathbb{J}, \mathbb{R}) \rightarrow C_{rd}(\mathbb{J}, \mathbb{R})$ defined by

$$S(x)(t) := E_p(b-a)\eta\phi(t)\varepsilon + L_f \int_a^t e_{\ominus p}(t, s)x(s)\Delta s. \tag{3.7}$$

For $u, v \in C_{rd}(\mathbb{J}, \mathbb{R}^+)$, we can write

$$S(u)(t) - S(v)(t) = L_f \int_a^t e_{\ominus p}(t, s)(u(s) - v(s))\Delta s.$$

Then

$$|S(u)(t) - S(v)(t)| \leq L_f E_p(b-a)\|u - v\|.$$

Since $E_p L_f (b-a) < 1$, we obtain that S is a contraction on $C_{rd}(\mathbb{J}, \mathbb{R})$, and using the Banach contraction principle, we see that S is a Picard operator and $F_S = \{u^*\}$. Then for $t \in \mathbb{J}$, we have

$$u^*(t) = E_p(b-a)\eta\phi(t)\varepsilon + L_f \int_a^t e_{\ominus p}(t, s)u^*(s)\Delta s.$$

We notice that u^* is increasing and

$$u^*(t) \leq E_p(b-a)\eta\phi(t)\varepsilon + \int_a^t E_p L_f u^*(s)\Delta s.$$

Employing the Gronwall inequality given in Theorem 2.1 to the above inequality with $y(t) = u^*(t)$, $\mathcal{F}(t) = E_p(b-a)\eta\phi(t)\varepsilon$, $\mathcal{H}(t) = E_p L_f$, and $G(t) = 0$, we obtain

$$u^*(t) \leq E_p(b-a)\eta\phi(t)\varepsilon e_{E_p L_f}(t, a) \quad \text{for all } t \in \mathbb{J}.$$

**Certified as
TRUE COPY**


Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

From (3.6), we have $u(t) \leq S(u)(t)$ for all $t \in \mathbb{J}$, where $u(t) = |y(t) - x(t)|$. Thus, S is an increasing Picard operator on $C_{rd}(\mathbb{J}, \mathbb{R})$. Now, in view of Lemma 2.1, we obtain $u(t) \leq u^*(t)$ for all $t \in \mathbb{J}$. This implies that

$$u(t) \leq E_p(b-a)e_{E_p L_f}(t, a)\varepsilon\eta\phi(t) \quad \text{for all } t \in \mathbb{J}.$$

That is,

$$|y(t) - x(t)| \leq E_p(b-a)e_{E_p L_f}(t, a)\varepsilon\eta\phi(t) \quad \text{for all } t \in \mathbb{J}.$$

Thus, the NDE (1.1) has Hyers–Ulam–Rassias stability of type \mathcal{N} with HURS constant $E_p(b-a)\eta e_{E_p L_f}(b, a)$. \square

Corollary 3.1. *Assume (C₁)–(C₃). If $E_p L_f(b-a) < 1$, then (1.1) has generalized Hyers–Ulam–Rassias stability of type \mathcal{N} with GHURS constant $E_p(b-a)\eta e_{E_p L_f}(b, a)$.*

Proof. In the proof of Theorem 3.1, if we take $\varepsilon = 1$, then we obtain

$$|y(t) - x(t)| \leq E_p(b-a)e_{E_p L_f}(b, a)\eta\phi(t) \quad \text{for all } t \in \mathbb{J}.$$

This shows that NDE (1.1) has generalized Hyers–Ulam–Rassias stability of type \mathcal{N} with GHURS constant $E_p(b-a)\eta e_{E_p L_f}(b, a)$. \square

Corollary 3.2. *Assume (C₁)–(C₃). If $E_p L_f(b-a) < 1$, then NDE (1.1) has Hyers–Ulam stability with HUS constant $E_p(b-a)\eta e_{E_p L_f}(b, a)$.*

Proof. In the proof of Theorem 3.1, if we take $\phi(t) \equiv 1$, then we obtain

$$|y(t) - x(t)| \leq E_p(b-a)\eta e_{E_p L_f}(b, a)\varepsilon \quad \text{for all } t \in \mathbb{J}.$$

Thus NDE (1.1) has Hyers–Ulam stability with HUS constant $E_p(b-a)\eta e_{E_p L_f}(b, a)$. \square

Corollary 3.3. *Assume (C₁)–(C₃). If $E_p L_f(b-a) < 1$, then NDE (1.1) has generalized Hyers–Ulam stability.*

Proof. Using $\theta_f(\varepsilon) = E_p(b-a)\eta e_{E_p L_f}(b, a)\varepsilon$, the result follows from Corollary 3.2. \square

Now, as an application of Theorem 3.1, we shall discuss Hyers–Ulam–Rassias stability of the adjoint equation to (1.1), namely

$$x^\Delta(t) + q(t)x(t) = g(t, x(t)) \quad \text{for all } t \in \mathbb{J}^K, \quad (3.8)$$

where $q \in \mathcal{R}^+(\mathbb{J}, \mathbb{R})$ and $g \in C_{rd}(\mathbb{J} \times \mathbb{R}, \mathbb{R})$.

Theorem 3.2. *Consider the adjoint NDE (3.8). Assume that the following conditions are satisfied.*

- (C₄) *Let $g \in C_{rd}(\mathbb{J} \times \mathbb{R}, \mathbb{R})$ and $q \in \mathcal{R}^+(\mathbb{J}, \mathbb{R})$ be such that $1 - \mu(t)q(t) > 0$ for all $t \in \mathbb{J}$.*
- (C₅) *There exists $L_g > 0$ such that*

**Certified as
TRUE COPY**


Principal

Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.

$$|g(t, u) - g(t, v)| \leq L_g(1 - \mu(t)q(t))|u - v| \quad (3.9)$$

for all $t \in \mathbb{J}$ and $u, v \in \mathbb{R}$.

(C₆) There exists $\eta > 0$ such that for $\phi \in \mathcal{N}(\mathbb{J}, \mathbb{R}^+)$

$$\int_a^t \phi(s) \Delta s \leq \eta \phi(t) \quad \text{for all } t \in \mathbb{J}. \quad (3.10)$$

If $E_q L_g(1 - \mu(t)q(t))(b - a) < 1$, then the following assertions hold:

- (i) The NDE (3.8) has a unique solution $x \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ satisfying the initial condition $x(a) = A$ for any initial value $A \in \mathbb{R}$.
- (ii) The NDE (3.8) has Hyers–Ulam–Rassias stability of type \mathcal{N} with HURS constant $E_q(b - a)\eta e_{E_q L_g}(b, a)$.

Proof. In view of the ‘simple useful formula’, putting $x^\sigma - \mu x^\Delta$ in place of x in (3.8), we obtain

$$x^\Delta(t) + q(t)(x^\sigma(t) - \mu(t)x^\Delta(t)) = g(t, x(t)).$$

Rearranging the terms, we can write this equation as

$$x^\Delta(t) + \left(\frac{q(t)}{1 - \mu(t)q(t)} \right) x^\sigma(t) = \frac{g(t, x(t))}{1 - \mu(t)q(t)}.$$

This equation is in the form of (1.1) with $p(t) := \frac{q(t)}{1 - \mu(t)q(t)}$ and $f(t, x) := \frac{g(t, x)}{1 - \mu(t)q(t)}$. In view of conditions (C₄) and (C₅), it is not difficult to show that (C₂) and (C₃) are verified. Hence, assertions (i) and (ii) follow from Theorem 3.1. \square

Remark 3.1. Other Ulam stability results for NDE (3.8) can be derived by using Theorem 3.2.

Another application of Theorem 3.1 concerns Hyers–Ulam–Rassias stability of the NDE

$$x^\Delta(t) = F(t, x(t)) \quad \text{for all } t \in \mathbb{J}^\kappa. \quad (3.11)$$

Theorem 3.3. Consider the NDE (3.11). Assume that the following conditions are satisfied.

(C₇) Let $F \in C_{rd}(\mathbb{J} \times \mathbb{R}, \mathbb{R})$ and $q \in \mathcal{R}^+(\mathbb{J}, \mathbb{R})$ be such that

$$1 - \mu(t)q(t) > |q(t)| \quad \text{for all } t \in \mathbb{J}.$$

(C₈) There exists $L_F > \frac{|q(t)|}{1 - \mu(t)q(t)}$ such that

$$|F(t, u) - F(t, v)| \leq (L_F(1 - \mu(t)q(t)) - |q(t)|)|u - v|$$

for all $t \in \mathbb{J}$ and $u, v \in \mathbb{R}$, where q is as given in (C₇).

(C₉) There exists $\eta > 0$ such that for $\phi \in \mathcal{N}(\mathbb{J}, \mathbb{R}^+)$

$$\int_a^t \phi(s) \Delta s \leq \eta \phi(t) \quad \text{for all } t \in \mathbb{J}. \quad (3.12)$$

If $(E_q L_F(1 - \mu(t)q(t)) - |q(t)|)(b - a) < 1$, then the following assertions hold:

**Certified as
TRUE COPY**


Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

- (i) The NDE (3.11) has a unique solution $x \in C_{\text{rd}}^1(\mathbb{J}, \mathbb{R})$ satisfying the initial condition $x(a) = A$ for any initial value $A \in \mathbb{R}$.
- (ii) The NDE (3.11) has Hyers–Ulam–Rassias stability of type \mathcal{N} with HURS constant $E_q(b-a)\eta e_{E_q L_F}(b, a)$.

Proof. Keeping in mind the ‘simple useful formula’ and rearranging the terms, we rewrite (3.11) as

$$x^\Delta(t) + \left(\frac{q(t)}{1 - \mu(t)q(t)} \right) x^\sigma(t) = \frac{q(t)x(t) + F(t, x(t))}{1 - \mu(t)q(t)} \quad \text{for all } t \in \mathbb{J}^\kappa.$$

That is,

$$x^\Delta(t) + p(t)x^\sigma(t) = f(t, x(t)) \quad \text{for all } t \in \mathbb{J}^\kappa,$$

where $p(t) := \frac{q(t)}{1 - \mu(t)q(t)}$ and $f(t, x) := \frac{q(t)x + F(t, x)}{1 - \mu(t)q(t)}$. Now, it remains to verify the condition (C₂) in Theorem 3.1. From (C₈), we have

$$|F(t, u) - F(t, v)| + |q(t)||u - v| \leq L_F(1 - \mu(t)q(t))|u - v|$$

for all $t \in \mathbb{J}$ and $u, v \in \mathbb{R}$. This gives

$$|F(t, u) - F(t, v) + q(t)(u - v)| \leq L_F(1 - \mu(t)q(t))|u - v|$$

for all $t \in \mathbb{J}$ and $u, v \in \mathbb{R}$. Rearranging the terms, we can write

$$\left| \frac{F(t, u) + q(t)u}{1 - \mu(t)q(t)} - \frac{F(t, v) + q(t)v}{1 - \mu(t)q(t)} \right| \leq L_F|u - v|$$

for all $t \in \mathbb{J}$ and $u, v \in \mathbb{R}$. That is,

$$|f(t, u) - f(t, v)| \leq L_F|u - v| \quad \text{for all } t \in \mathbb{J} \text{ and } u, v \in \mathbb{R}.$$

Hence, (C₂) is verified. Now, we are able to apply Theorem 3.1 and obtain that assertions (i) and (ii) follow from Theorem 3.1. \square

Remark 3.2. Other Ulam stability results for NDE (3.11) can be derived by using Theorem 3.3.

4. EXAMPLE

Let $\mathbb{T} = 2^{\mathbb{N}_0}$ and $a = 2, b = 32$. Then $\mathbb{J} := [a, b] \cap \mathbb{T} = \{2, 4, 8, 16, 32\}$. Consider the NDE

$$x^\Delta(t) + tx^\sigma(t) = \frac{1}{e_t(t, 2)} + \frac{(x^2(t) + 5)^{1/2}}{8} \quad \text{for all } t \in \mathbb{J}^\kappa \quad (4.1)$$

with the initial condition $x(2) = 2$ and the inequality

$$\left| y^\Delta(t) + ty^\sigma(t) - \frac{1}{e_t(t, 2)} - \frac{(y^2(t) + 5)^{1/2}}{8} \right| \leq \varepsilon e_{\frac{1}{20}}(t, 2) \quad \text{for all } t \in \mathbb{J}^\kappa. \quad (4.2)$$

Here $f(t, x(t)) = \frac{1}{e_t(t, 2)} + \frac{(x^2(t) + 5)^{1/2}}{8}$ which satisfies (C₂) with $L_f = \frac{1}{8}$, and $p = t$, $\mu(t) = t$ for all $t \in \mathbb{T}$. Clearly, $1 + \mu(t)p(t) = 1 + t^2 > 0$. Thus $p \in \mathcal{R}^+(\mathbb{J}, \mathbb{R})$. Also,

Certified as
TRUE COPY



Principal

Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.

$$(\ominus p)(t) = \frac{-p(t)}{1 + \mu(t)p(t)} = \frac{-t}{1 + t^2}.$$

With these values, we obtain

$$\begin{aligned} e_{\ominus p}(t, s) &= \exp \left(\int_s^t \frac{1}{\mu(r)} \ln(1 + \mu(r) \ominus p) \Delta r \right) \\ &= \exp \left(\int_s^t \frac{1}{r} \ln \left(1 + r \left(\frac{-r}{1 + r^2} \right) \right) \Delta r \right) \\ &= \exp \left(\int_s^t \frac{1}{r} \ln \left(1 + \frac{-r^2}{1 + r^2} \right) \Delta r \right) \\ &= \exp \left(\int_s^t \frac{1}{r} \ln \left(\frac{1}{1 + r^2} \right) \Delta r \right) \\ &= \exp \left(\sum_{r=s}^{t/2} \ln \left(\frac{1}{1 + r^2} \right) \right) \\ &= \prod_{r=s}^{t/2} \frac{1}{1 + r^2} \quad \text{for } s \leq t \text{ and } t \in \mathbb{J}. \end{aligned}$$

This leads to

$$E_p = \sup_{s, t \in \mathbb{J}} \left| \prod_{r=s}^{t/2} \frac{1}{1 + r^2} \right| = \frac{1}{5}.$$

Further, we find that $E_p L_f(b-a) = \frac{1}{5} \frac{1}{8} (32-2) = \frac{3}{4} < 1$. Thus, all the conditions in Theorem 3.1 are satisfied. Therefore, (4.1) has a unique solution satisfying initial condition $x(2) = 2$. Now, let $y \in C_{rd}^1(\mathbb{J}, \mathbb{R})$ be a solution of (4.2). Then, by Remark 2.3, there exists $\psi \in C_{rd}(\mathbb{J}, \mathbb{R})$ such that $|\psi(t)| \leq \epsilon e_{\frac{1}{20}}(t, 2)$ and

$$y^\Delta(t) + ty^\sigma(t) = \frac{1}{e_p(t, 2)} + \frac{(y^2(t) + 5)^{1/2}}{8} + \psi(t) \quad \text{for all } t \in \mathbb{J}^k. \quad (4.3)$$

By Lemma 2.2, we have

$$\begin{aligned} y(t) &= (y(2) + t - 2) \prod_{r=2}^{t/2} \frac{1}{1 + r^2} + \frac{1}{8} \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1 + r^2} \right) (y^2(r) + 5)^{1/2} \\ &\quad + \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1 + r^2} \right) \psi(s) \quad \text{for all } t \in \mathbb{J}. \end{aligned} \quad (4.4)$$

From Theorem 3.1, we find that the dynamic problem

$$\begin{aligned} x^\Delta(t) + tx^\sigma(t) &= \frac{1}{e_p(t, 2)} + \frac{(x^2(t) + 5)^{1/2}}{8}; \quad t \in \mathbb{J}^k, \\ x(2) &= y(2) \end{aligned}$$

has a unique solution. According to Lemma 2.2, this unique solution is given by

**Certified as
TRUE COPY**



Principal

**Ramniranjan Jhunjunwala College,
Ghatkopar (W), Mumbai-400086.**

$$\begin{aligned}
x(t) &= y(2)e_{\ominus p}(t, 2) + \int_2^t e_{\ominus p}(t, s) \left(\frac{1}{e_p(s, 2)} + \frac{(x^2(s) + 5)^{1/2}}{8} \right) \Delta s \\
&= y(2)e_{\ominus p}(t, 2) + \int_2^t \left(e_{\ominus p}(t, 2) + e_{\ominus p}(t, s) \frac{(x^2(s) + 5)^{1/2}}{8} \right) \Delta s \\
&= y(2)e_{\ominus p}(t, 2) + e_{\ominus p}(t, 2)(t - 2) + \frac{1}{8} \int_2^t e_{\ominus p}(t, s) (x^2(s) + 5)^{1/2} \Delta s \\
&= (y(2) + t - 2)e_{\ominus p}(t, 2) + \frac{1}{8} \int_2^t e_{\ominus p}(t, s) (x^2(s) + 5)^{1/2} \Delta s \\
&= (y(2) + t - 2) \prod_{r=2}^{t/2} \frac{1}{1+r^2} + \frac{1}{8} \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1+r^2} \right) (x^2(r) + 5)^{1/2}.
\end{aligned}$$

That is,

$$\begin{aligned}
x(t) &= (y(2) + t - 2) \prod_{r=2}^{t/2} \frac{1}{1+r^2} \\
&\quad + \frac{1}{8} \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1+r^2} \right) (x^2(r) + 5)^{1/2} \quad \text{for all } t \in \mathbb{J}.
\end{aligned} \tag{4.5}$$

Now, from (4.4) and (4.5), we can write for $t \in \mathbb{J}$,

$$\begin{aligned}
|y(t) - x(t)| &\leq \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1+r^2} \right) |\Psi(s)| \\
&\leq \sum_{s=2}^{t/2} \left(\prod_{r=s}^{t/2} \frac{1}{1+r^2} \right) \varepsilon e_{\frac{1}{20}}(s, 2) \\
&\leq \varepsilon \frac{1}{5} \sum_{s=2}^{t/2} e_{\frac{1}{20}}(s, 2) \\
&\leq \varepsilon \frac{1}{5} e_{\frac{1}{20}}(32, 2) 30.
\end{aligned}$$

Thus, $|y(t) - x(t)| \leq 6\varepsilon e_{\frac{1}{20}}(32, 2)$, $t \in \mathbb{J}$, which yields that (4.1) has Hyers–Ulam–Rassias stability of type \mathcal{N} with HURS constant $6e_{\frac{1}{20}}(32, 2)$.

ACKNOWLEDGEMENTS

The authors thank both referees for carefully reading this manuscript and making many important suggestions, leading to a better presentation of the results.

REFERENCES

- [1] Maryam A. Alghamdi, Mymonah Alharbi, Martin Bohner, and Alaa E. Hamza. *Hyers–Ulam and Hyers–Ulam–Rassias stability of first-order nonlinear dynamic equations*. Qual. Theory Dyn. Syst., 20(2):Paper No. 45, 14, 2021.

**Certified as
TRUE COPY**

Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

- [2] Maryam A. Alghamdi, Alaa Aljehani, Martin Bohner, and Alaa E. Hamza. *Hyers–Ulam and Hyers–Ulam–Rassias stability of first-order linear dynamic equations*. Publ. Inst. Math. (Beograd) (N.S.), 109(123):83–93, 2021.
- [3] Qusuay H. Alqifiary and Soon-Mo Jung. *On the Hyers–Ulam stability of differential equations of second order*. Abstr. Appl. Anal., Art. ID 483707, 8 pages, 2014.
- [4] Douglas R. Anderson and Masakazu Onitsuka. *Hyers–Ulam stability of first-order homogeneous linear dynamic equations on time scales*. Demonstr. Math., 51(1):198–210, 2018.
- [5] Douglas R. Anderson and Masakazu Onitsuka. *Hyers–Ulam stability for a discrete time scale with two step sizes*. Appl. Math. Comput., 344/345:128–140, 2019.
- [6] Douglas R. Anderson and Masakazu Onitsuka. *Hyers–Ulam stability for Cayley quantum equations and its application to h -difference equations*. Mediterr. J. Math., 18(4):Paper No.168, 13, 2021.
- [7] Douglas R. Anderson and Masakazu Onitsuka. *Hyers–Ulam stability for quantum equations*. Aequationes Math., 95(2):201–214, 2021.
- [8] Douglas R. Anderson and Masakazu Onitsuka. *Ulam stability for nonautonomous quantum equations*. J. Inequal. Appl., Paper No. 161, 16 pages, 2021.
- [9] Szilárd András and Alpár Richárd Mészáros. *Ulam–Hyers stability of dynamic equations on time scales via Picard operators*. Appl. Math. Comput., 219(9):4853–4864, 2013.
- [10] Tosio Aoki. *On the stability of the linear transformation in Banach spaces*. J. Math. Soc. Japan, 2:64–66, 1950.
- [11] Alina Ramona Baias, Florina Blaga, and Dorian Popa. *Best Ulam constant for a linear difference equation*. Carpathian J. Math., 35(1):13–22, 2019.
- [12] Alina Ramona Baias and Dorian Popa. *On Ulam stability of a linear difference equation in Banach spaces*. Bull. Malays. Math. Sci. Soc., 43(2):1357–1371, 2020.
- [13] Alina Ramona Baias and Dorian Popa. *On the best Ulam constant of a higher order linear difference equation*. Bull. Sci. Math., 166:Paper No. 102928, 12, 2021.
- [14] Abdellatif Ben Makhlouf, Lassaad Mchiri, and Mohamed Rhaima. *Ulam–Hyers–Rassias stability of stochastic functional differential equations via fixed point methods*. J. Funct. Spaces, Art. ID 5544847, 7 pages, 2021.
- [15] Arzu Bilgin and Mustafa R. S. Kulenović. *Global asymptotic stability for discrete single species population models*. Discrete Dyn. Nat. Soc., Art. ID 5963594, 15 pages, 2017.
- [16] Martin Bohner and Allan Peterson. *Dynamic equations on time scales*. Birkhäuser Boston, Inc., Boston, MA, 2001. An introduction with applications.
- [17] Martin Bohner and Allan Peterson. *Advances in dynamic equations on time scales*. Birkhäuser Boston, Inc., Boston, MA, 2003.
- [18] Martin Bohner, Sanket Tikare, and Iguer Luis Domini dos Santos. *First-order nonlinear dynamic initial value problems*. Int. J. Dyn. Syst. Differ. Equ., 11(3-4):241–254, 2021.
- [19] David G. Bourgin. *Classes of transformations and bordering transformations*. Bull. Amer. Math. Soc., 57:223–237, 1951.
- [20] Vahidin Hadžiabdić, Mustafa R. S. Kulenović, and Esmir Pilav. *Global stability of a quadratic anti-competitive system of rational difference equations in the plane with Allee effects*. J. Comput. Anal. Appl., 25(6):1132–1144, 2018.
- [21] Jinghao Huang, Soon-Mo Jung, and Yongjin Li. *On Hyers–Ulam stability of nonlinear differential equations*. Bull. Korean Math. Soc., 52(2):685–697, 2015.
- [22] Donald H. Hyers. *On the stability of the linear functional equation*. Proc. Nat. Acad. Sci. U.S.A., 27:222–224, 1941.
- [23] Sabina Jašarević Hrustić, Mustafa R. S. Kulenović, and Mehmed Nurkanović. *Local dynamics and global stability of certain second-order rational difference equation with quadratic terms*. Discrete Dyn. Nat. Soc., Art. ID 3716042, 14 pages, 2016.

**Certified as
TRUE COPY**


Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

- [24] Mustafa R. S. Kulenović, Samra Moranjkić, Mehmed Nurkanović, and Zehra Nurkanović. *Global asymptotic stability and Naimark–Sacker bifurcation of certain mix monotone difference equation*. Discrete Dyn. Nat. Soc., Art. ID 7052935, 22 pages, 2018.
- [25] Tongxing Li, Nicola Pintus, and Giuseppe Viglialoro. *Properties of solutions to porous medium problems with different sources and boundary conditions*. Z. Angew. Math. Phys., 70(3):Paper No. 86, 18 pages, 2019.
- [26] Tongxing Li and Giuseppe Viglialoro. *Boundedness for a nonlocal reaction chemotaxis model even in the attraction-dominated regime*. Differential Integral Equations, 34(5-6):315–336, 2021.
- [27] Adela Novac, Diana Otrocol, and Dorian Popa. *Ulam stability of a linear difference equation in locally convex spaces*. Results Math., 76(1):Paper No. 33, 13, 2021.
- [28] Masakazu Onitsuka. *Influence of the stepsize on Hyers–Ulam stability of first-order homogeneous linear difference equations*. Int. J. Difference Equ., 12(2):281–302, 2017.
- [29] Masakazu Onitsuka and Tomohiro Shoji. *Hyers–Ulam stability of first-order homogeneous linear differential equations with a real-valued coefficient*. Appl. Math. Lett., 63:102–108, 2017.
- [30] Dorian Popa. *Hyers–Ulam stability of the linear recurrence with constant coefficients*. Adv. Difference Equ., (2):101–107, 2005.
- [31] Themistocles M. Rassias. *On the stability of the linear mapping in Banach spaces*. Proc. Amer. Math. Soc., 72(2):297–300, 1978.
- [32] Ioan A. Rus. *Gronwall lemmas: ten open problems*. Sci. Math. Jpn., 70(2):221–228, 2009.
- [33] Ioan A. Rus. *Ulam stability of ordinary differential equations*. Stud. Univ. Babeş-Bolyai Math., 54(4):125–133, 2009.
- [34] Ioan A. Rus. *Ulam stabilities of ordinary differential equations in a Banach space*. Carpathian J. Math., 26(1):103–107, 2010.
- [35] Yonghong Shen. *The Ulam stability of first order linear dynamic equations on time scales*. Results Math., 72(4):1881–1895, 2017.
- [36] Yonghong Shen and Yongjin Li. *A general method for the Ulam stability of linear differential equations*. Bull. Malays. Math. Sci. Soc., 42(6):3187–3211, 2019.
- [37] Sanket Tikare. *Nonlocal initial value problems for first-order dynamic equations on time scales*. Appl. Math. E-Notes, 21:410–420, 2021.
- [38] Sanket Tikare, Martin Bohner, Bipan Hazarika, and Ravi P Agarwal. *Dynamic local and nonlocal initial value problems in Banach spaces*. Rend.Circ.Mat.Palermo, II. Ser (2021), 1–16, 2021.
- [39] Stanislaw M. Ulam. *A collection of mathematical problems*. Interscience Tracts in Pure and Applied Mathematics, no. 8. Interscience Publishers, New York-London, 1960.
- [40] Fu-Hsiang Wong, Cheh-Chih Yeh, and Chen-Huang Hong. *Gronwall inequalities on time scales*. Math. Inequal. Appl., 9(1):75–86, 2006.

(Received: February 20, 2022)

(Revised: April 01, 2022)

Martin Bohner
Missouri S&T
Department of Mathematics and Statistics
Rolla, MO 65409-0020, USA.
e-mail: bohner@mst.edu

and

Sanket Tikare
Ramniranjan Jhunjhunwala College
Department of Mathematics
Mumbai, Maharashtra 400 086, India.
e-mail: sankettikare@rjcollege.edu.in

**Certified as
TRUE COPY**



Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**

41. Zada, A., Shah, S.O.: Hyers-Ulam stability of first-order non-linear delay differential equations with fractional integrable impulses. *Hacet. J. Math. Stat.* **47**(5), 1196–1205 (2018)
42. Zhu, Z.-Q., Wang, Q.-R.: Existence of nonoscillatory solutions to neutral dynamic equations on time scales. *J. Math. Anal. Appl.* **335**(2), 751–762 (2007)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

**Certified as
TRUE COPY**



Principal

**Ramniranjan Jhunjhunwala College,
Ghatkopar (W), Mumbai-400086.**