



Scopus Preview



## Source details

### Applied Mathematics E - Notes

Scopus coverage years: from 2001 to 2021

Publisher: Tsinghua University

ISSN: 1607-2510

Subject area: Mathematics: Applied Mathematics

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

CiteScore 2020

0.5

SJR 2020

0.241

SNIP 2020

0.520

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

#### Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020 ▼

$$0.5 = \frac{96 \text{ Citations 2017 - 2020}}{177 \text{ Documents 2017 - 2020}}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021 ⓘ

$$0.8 = \frac{177 \text{ Citations to date}}{220 \text{ Documents to date}}$$

Last updated on 06 April, 2022 • Updated monthly

#### CiteScore rank 2020 ⓘ

Category	Rank	Percentile
Mathematics		
└ Applied Mathematics	#477/548	12th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

**Certified as  
TRUE COPY**

**Principal**

**Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.**

ISSN 1607-2510

HOME

ALL VOLUMES

**Editor-in-Chief**

Sui-Sun Cheng

Address

**Editorial Board**

S. S. Cheng

M. Gil'

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

List

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

Address

**Technical Advisor**

Y. C. Chang

# Applied Mathematics E-Notes

Published Contents of Volume 21

1. Mohamad A. Fawzy, Haseeb Athar and Yousef F. Alharbi
  - Inference based on Marshall-Olkin extended Rayleigh Lomax distribution
  - *Appl. Math. E-Notes*, 21(2021), 1-11
  - Retrieve article in: [PDF](#)
2. Hüseyin Budak, Candan Can Bilişik, Artion Kashuri and Muhammad Aamir Ali
  - Hermite-Hadamard type inequalities for the interval-valued harmonically h-convex functions via fractional integrals
  - *Appl. Math. E-Notes*, 21(2021), 12-32
  - Retrieve article in: [PDF](#)
3. Robert Frontczak
  - A remark on the crux problem 1052\*
  - *Appl. Math. E-Notes*, 21(2021), 33-36
  - Retrieve article in: [PDF](#)
4. Horst Alzer and Rui A. C. Ferreira
  - Concavity and generalized entropy
  - *Appl. Math. E-Notes*, 21(2021), 37-43
  - Retrieve article in: [PDF](#)
5. Mohammed Massar
  - On a nonlocal Schrödinger-Poisson system with critical exponent
  - *Appl. Math. E-Notes*, 21(2021), 44-52
  - Retrieve article in: [PDF](#)
6. Sharafat Hussain
  - Fixed point theorems for nonlinear contraction in controlled metric type space
  - *Appl. Math. E-Notes*, 21(2021), 53-61
  - Retrieve article in: [PDF](#)
7. Sevda Yıldız
  - $K_\alpha$ -convergence for double sequences and Korovkin type approximation
  - *Appl. Math. E-Notes*, 21(2021), 62-71
  - Retrieve article in: [PDF](#)
8. Kuppathai Appasamy Selvakumaran, Junesang Choi and Sunil Dutt Purohit
  - Certain subclasses of analytic functions defined by fractional q-calculus operators
  - *Appl. Math. E-Notes*, 21(2021), 72-80
  - Retrieve article in: [PDF](#)
9. Ekin Uğurlu and Kenan Tas
  - Dependence of eigenvalues of some boundary value problems
  - *Appl. Math. E-Notes*, 21(2021), 81-88
  - Retrieve article in: [PDF](#)
10. Ioannis Konstantinos Argyros and Santhosh George
  - Extended convergence of Jarratt type methods
  - *Appl. Math. E-Notes*, 21(2021), 89-96
  - Retrieve article in: [PDF](#)
11. İbrahim Adalar
  - A uniqueness theorem for inverse Sturm-Liouville problem
  - *Appl. Math. E-Notes*, 21(2021), 97-102

Certified as TRUE COPY

Principal

Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



ISSN 1607-2510

HOME

ALL VOLUMES

**Editor-in-Chief**

Sui-Sun Cheng

Address

**Editorial Board**

S. S. Cheng

M. Gil'

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

List

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

Address

**Technical Advisor**

Y. C. Chang

12. Kadri Arslan and Betül Bulca

- Rotational surfaces with rotations in  $x_3x_4$ -plane
- *Appl. Math. E-Notes*, 21(2021), 103-109
- Retrieve article in: [PDF](#)

13. Gamaliel Cerda-Morales

- Investigation of generalized Fibonacci hybrid numbers and their properties
- *Appl. Math. E-Notes*, 21(2021), 110-118
- Retrieve article in: [PDF](#)

14. Mohamed Grazem, Ahmed Bendjeddou and Rachid Cheurfa

- Systems having Liouvillian first integrals and non-algebraic limit cycles
- *Appl. Math. E-Notes*, 21(2021), 119-128
- Retrieve article in: [PDF](#)

15. Mohamed Akkouchi and Mohamed Amine Ighachane

- A generalized refinement of Young Inequality and applications
- *Appl. Math. E-Notes*, 21(2021), 129-138
- Retrieve article in: [PDF](#)

16. Shyam S. Kamath, A. Senthil Thilak and Rashmi Maladi

- Bounds on k-part degree restricted domination number of a graph
- *Appl. Math. E-Notes*, 21(2021), 139-151
- Retrieve article in: [PDF](#)

17. Indrajit Lahiri and Shubhashish Das

- A note on a conjecture of R. Brück
- *Appl. Math. E-Notes*, 21(2021), 152-156
- Retrieve article in: [PDF](#)

18. Tayeb Salhi

- On a certain class of Kolmogorov systems: integrability and non-existence of limit cycles
- *Appl. Math. E-Notes*, 21(2021), 157-162
- Retrieve article in: [PDF](#)

19. Davod Khojasteh Salkuyeh

- On computing sparse approximate inverse factor of SPD matrices
- *Appl. Math. E-Notes*, 21(2021), 163-171
- Retrieve article in: [PDF](#)

20. Saudia Jabeen, Muhammad Aslam Noor and Khalida Inayat Noor

- Some new inertial projection methods for Quasi variational inequalities
- *Appl. Math. E-Notes*, 21(2021), 172-178
- Retrieve article in: [PDF](#)

21. Zahra Goodarzi, Mohammad Reza Mokhtarzadeh, Mohammad Reza Pournaki and Abdolrahman Razani

- A note on periodic solutions of matrix Riccati differential equations
- *Appl. Math. E-Notes*, 21(2021), 179-186
- Retrieve article in: [PDF](#)

22. Mehdi Hassani

- On the asymptotic expansion of a generalized Smith's determinant
- *Appl. Math. E-Notes*, 21(2021), 187-193
- Retrieve article in: [PDF](#)

23. Sayyed Hashem Rasouli

- A note on the stability of nonnegative solutions to classes of fractional Laplacian problems with concave nonlinearity
- *Appl. Math. E-Notes*, 21(2021), 194-197
- Retrieve article in: [PDF](#)

**Certified as  
TRUE COPY**



**Principal**  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



ISSN 1607-2510

HOME

ALL VOLUMES

**Editor-in-Chief**

Sui-Sun Cheng

Address

**Editorial Board**

S. S. Cheng

M. Gil'

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

List

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

Address

**Technical Advisor**

Y. C. Chang

24. Saqib Mubarak, Mukhtar Ahmad Khanday, Ahsan Ul Haq Lone and Namreen Rasool

- An analytical approach to study the drug diffusion through transdermal drug delivery system
- *Appl. Math. E-Notes*, 21(2021), 198-208
- Retrieve article in: [PDF](#)

25. Kushal Roy, Sayantan Panja and Mantu Saha

- A generalized fixed point theorem in an extended cone  $b$ -metric space over Banach algebra with its application to dynamical programming
- *Appl. Math. E-Notes*, 21(2021), 209-219
- Retrieve article in: [PDF](#)

26. Luis Henri Gallardo

- On Ryser's conjecture: modulo 2 approach
- *Appl. Math. E-Notes*, 21(2021), 220-224
- Retrieve article in: [PDF](#)

27. Meena Joshi, Anita Tomar and Sanjay Kumar Padaliya

- Fixed point to fixed ellipse in Metric spaces and discontinuous activation function
- *Appl. Math. E-Notes*, 21(2021), 225-237
- Retrieve article in: [PDF](#)

28. Ahmed Chaouki Aouine and Abdelkrim Aliouche

- Fixed point theorems of Kannan type with an application to control theory
- *Appl. Math. E-Notes*, 21(2021), 238-249
- Retrieve article in: [PDF](#)

29. Sattar Alyar and Rana Khoeilar

- On the sum-connectivity index of trees
- *Appl. Math. E-Notes*, 21(2021), 250-252
- Retrieve article in: [PDF](#)

30. Nurten Kılıç, Özkan Öcalan and Umut Mutlu Özkan

- Oscillation tests for nonlinear differential equations with several nonmonotone advanced arguments
- *Appl. Math. E-Notes*, 21(2021), 253-262
- Retrieve article in: [PDF](#)

31. Harendra Verma, Vishnu Narayan Mishra and Pankaj Mathur

- Analysis of the impact of media in spreading the blue whale game
- *Appl. Math. E-Notes*, 21(2021), 263-274
- Retrieve article in: [PDF](#)

32. Zaid Laadjal and Nacer Adjeroud

- Sharp estimates for the unique solution of the Hadamard-type two-point fractional boundary value problems
- *Appl. Math. E-Notes*, 21(2021), 275-281
- Retrieve article in: [PDF](#)

33. Riccardo Fazio

- Perpetual American put option: an error estimator for a second order non-standard finite difference scheme
- *Appl. Math. E-Notes*, 21(2021), 282-290
- Retrieve article in: [PDF](#)

34. Yacine Arioua and Maria Titrœui

- Boundary value problem for a coupled system of nonlinear fractional differential equations involving Erdélyi-Kober derivative
- *Appl. Math. E-Notes*, 21(2021), 291-306
- Retrieve article in: [PDF](#)

35. Anis Hezzam and Abderrahmane Turki

**Certified as  
TRUE COPY**



**Principal**  
Ramniranjan Jhunjunwala College,  
Ghatkopar (W), Mumbai-400086.



ISSN 1607-2510

HOME

ALL VOLUMES

**Editor-in-Chief**

Sui-Sun Cheng

Address

**Editorial Board**

S. S. Cheng

M. Gil

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

List

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

Address

**Technical Advisor**

Y. C. Chang

- Retrieve article in: [PDF](#)
- 35. Anis Hezzam and Abderrahmane Turki
  - On the affine equivalence and minimal rational bases of planar cubic differential systems
  - *Appl. Math. E-Notes*, 21(2021), 307-319
  - Retrieve article in: [PDF](#)
- 36. Parimala Sivakumar, Kalyanasundaram Madhu and Jayakumar Jayaraman
  - Optimal eighth and sixteenth order iterative methods for solving nonlinear equation with basins of attraction
  - *Appl. Math. E-Notes*, 21(2021), 320-343
  - Retrieve article in: [PDF](#)
- 37. Sarah Elahi, Muhammad Aslam Noor and Khalida Inayat Noor
  - Fractional integral inequalities for preinvex functions
  - *Appl. Math. E-Notes*, 21(2021), 344-355
  - Retrieve article in: [PDF](#)
- 38. S. Ithaya Ezhil Manna and A. Anthony Eldred
  - Iterative approximation of best proximity pairs of asymptotically relatively nonexpansive mappings
  - *Appl. Math. E-Notes*, 21(2021), 356-364
  - Retrieve article in: [PDF](#)
- 39. Salim Yüce
  - On the E. Study maps for the dual quaternions
  - *Appl. Math. E-Notes*, 21(2021), 365-374
  - Retrieve article in: [PDF](#)
- 40. Naila Mehreen and Matloob Anwar
  - Integral inequalities for exponentially  $(x_3x_4)$ -convex functions
  - *Appl. Math. E-Notes*, 21(2021), 375-384
  - Retrieve article in: [PDF](#)
- 41. Sirous Moradi, Ebrahim Analouei Audegani and Mohammad Mohammadi Anjedani
  - Existence and uniqueness of solutions of Volterra-Fredholm nonlinear integral equation in partially ordered  $b$ -metric spaces
  - *Appl. Math. E-Notes*, 21(2021), 385-401
  - Retrieve article in: [PDF](#)
- 42. Sai Gopal Rayaguru and Gopal Krishna Panda
  - Repdigits as sums of two associated Pell numbers
  - *Appl. Math. E-Notes*, 21(2021), 402-409
  - Retrieve article in: [PDF](#)
- 43. Sanket Tikare
  - Nonlocal initial value problems for first-order dynamic equations on time scales
  - *Appl. Math. E-Notes*, 21(2021), 410-420
  - Retrieve article in: [PDF](#)
- 44. Yogesh Jibhau Bagul
  - On the Shafer-type inequality for the inverse tangent function
  - *Appl. Math. E-Notes*, 21(2021), 421-425
  - Retrieve article in: [PDF](#)
- 45. Basem Aref Frasin
  - An application of generalized Bessel functions on general class of analytic functions with negative coefficients
  - *Appl. Math. E-Notes*, 21(2021), 426-436
  - Retrieve article in: [PDF](#)
- 46. Yousaf Khurshid and Muhammad Adil Khan
  - Hermite-Hadamard type inequalities for conformable integrals via

Principal

Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



ISSN 1607-2510

[HOME](#)

[ALL VOLUMES](#)

**Editor-in-Chief**

Sui-Sun Cheng

[Address](#)

**Editorial Board**

S. S. Cheng

M. Gil'

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

[List](#)

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

[Address](#)

**Technical Advisor**

Y. C. Chang

- Retrieve article in: [PDF](#)
- 45. Basem Aref Frasin
  - An application of generalized Bessel functions on general class of analytic functions with negative coefficients
  - *Appl. Math. E-Notes*, 21(2021), 426-436
  - Retrieve article in: [PDF](#)
- 46. Yousaf Khurshid and Muhammad Adil Khan
  - Hermite-Hadamard type inequalities for conformable integrals via preinvex functions
  - *Appl. Math. E-Notes*, 21(2021), 437-450
  - Retrieve article in: [PDF](#)
- 47. Jagdish Saran, Narinder Pushkarna and Kanika Verma
  - Progressively type-II right censored order statistics from odds generalized exponential-Pareto distribution and related inference
  - *Appl. Math. E-Notes*, 21(2021), 451-466
  - Retrieve article in: [PDF](#)
- 48. Anguraj Annamalai, Ravikumar Kasinathan, Ramkumar Kasinathan and Elsayed Mohammed Elsayed
  - Existence, uniqueness and stability results of impulsive stochastic semilinear neutral functional partial integrodifferential equations with infinite delay and Poisson jumps
  - *Appl. Math. E-Notes*, 21(2021), 467-477
  - Retrieve article in: [PDF](#)
- 49. Vineet Bhatt, Kailash Chandra Petwal, Sunil Kumar, Sakshi Rajput and Manoj
  - A comprehensive note on techniques used for face detection and application
  - *Appl. Math. E-Notes*, 21(2021), 478-494
  - Retrieve article in: [PDF](#)
- 50. Kapula Rajendra Prasad, Mahammad Khuddush and Pogadadanda Veeraiah
  - Denumerably many positive solutions for fractional differential equation with Riemann-Stieltjes integral boundary conditions
  - *Appl. Math. E-Notes*, 21(2021), 495-504
  - Retrieve article in: [PDF](#)
- 51. Bashir Ahmad, Ahmed Alsaedi, Mona Alsulami and Sotiris K. Ntouyas
  - On nonlinear third-order boundary value problems involving non-separated type strip-multi-point boundary conditions
  - *Appl. Math. E-Notes*, 21(2021), 505-524
  - Retrieve article in: [PDF](#)
- 52. Nisar Ahmad Rather, Ishfaq Dar and Mohammad Shafi
  - On the zero bounds of polynomials and related analytic functions
  - *Appl. Math. E-Notes*, 21(2021), 525-532
  - Retrieve article in: [PDF](#)
- 53. Ali Hakem
  - Non-existence of global solutions of semi-linear Moore-Gibson-Thompson equation on the Heisenberg group
  - *Appl. Math. E-Notes*, 21(2021), 533-541
  - Retrieve article in: [PDF](#)
- 54. Rahim Safshekan and Ali Farokhinia
  - Some refinements of numerical radius inequalities via convex functions
  - *Appl. Math. E-Notes*, 21(2021), 542-549
  - Retrieve article in: [PDF](#)
- 55. Ramkumar Kasinathan, Banupriya Kandhasamy and Varshini Sandrasekaran
  - Existence and stability results for impulsive differential evolution system

  
Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



ISSN 1607-2510

HOME

ALL VOLUMES

**Editor-in-Chief**

Sui-Sun Cheng

Address

**Editorial Board**

S. S. Cheng

M. Gil'

W. T. Li

Y. Z. Lin

T. Kusano

B. S. Tam

J. C. Yao

S. S. Dragomir

V. A. Galaktionov

V. Gupta

V. M. Ungureanu

A. Prykarpatsky

J. B. Li

R. Y. Ma

L. Sanchez

M. Aslam Noor

H. S. Ren

C. Cuevas

S. Y. Huang

**Referees**

List

**Past Editors**

X. D. Zhu

Z. C. Li

H. C. Lai

Z. X. Zheng

J. M. Almira

Yu. Domshlak

C. V. Pao

J. S. Yu

G. Zhang

T. K. Pogany

**Editorial Assistant**

Po Chieh Chen

Address

**Technical Advisor**

Y. C. Chang

- On nonlinear third-order boundary value problems involving non-separated type strip-multi-point boundary conditions
- *Appl. Math. E-Notes*, 21(2021), 505-524
- Retrieve article in: [PDF](#)
- 52. Nisar Ahmad Rather, Ishfaq Dar and Mohammad Shafi
  - On the zero bounds of polynomials and related analytic functions
  - *Appl. Math. E-Notes*, 21(2021), 525-532
  - Retrieve article in: [PDF](#)
- 53. Ali Hakem
  - Non-existence of global solutions of semi-linear Moore-Gibson-Thompson equation on the Heisenberg group
  - *Appl. Math. E-Notes*, 21(2021), 533-541
  - Retrieve article in: [PDF](#)
- 54. Rahim Safshekan and Ali Farokhnia
  - Some refinements of numerical radius inequalities via convex functions
  - *Appl. Math. E-Notes*, 21(2021), 542-549
  - Retrieve article in: [PDF](#)
- 55. Ramkumar Kasinathan, Banupriya Kandasamy and Varshini Sandrasekaran
  - Existence and stability results for integrodifferential evolution system with random impulse
  - *Appl. Math. E-Notes*, 21(2021), 550-562
  - Retrieve article in: [PDF](#)
- 56. Mohamed I. Abbas
  - On the coupled system of  $\diamond_r$ -Caputo fractional differential equations with four-point boundary conditions
  - *Appl. Math. E-Notes*, 21(2021), 563-576
  - Retrieve article in: [PDF](#)
- 57. Gaurav Dwivedi
  - Existence of multiple solutions for a Kirchhoff type equation involving polyharmonic operator with exponential growth
  - *Appl. Math. E-Notes*, 21(2021), 577-586
  - Retrieve article in: [PDF](#)
- 58. Kamel Tablennhas and Zoubir Dahmani
  - A three sequential fractional differential problem of Duffing type
  - *Appl. Math. E-Notes*, 21(2021), 587-598
  - Retrieve article in: [PDF](#)
- 59. Muhammad Ghaffar Khan, Bakhtiar Ahmad and Nak Eun Cho
  - Properties of Harmonic mappings associated with polylogarithm functions
  - *Appl. Math. E-Notes*, 21(2021), 599-609
  - Retrieve article in: [PDF](#)
- 60. Hyun-Woo Jin and Min-Young Lee
  - On characterizations of continuous distributions by independence property of the quotient-type k-th lower record value
  - *Appl. Math. E-Notes*, 21(2021), 610-614
  - Retrieve article in: [PDF](#)
- 61. Suzana Antunović and Damir Vukičević
  - Detecting communities under constraints in directed acyclic networks
  - *Appl. Math. E-Notes*, 21(2021), 615-621
  - Retrieve article in: [PDF](#)
- 62. Mesfin Mekuria Woldaregay and Gemechis File Duressa
  - Robust numerical scheme for solving singularly perturbed differential equations involving small delays
  - *Appl. Math. E-Notes*, 21(2021), 622-633

Certified as  
TRUE COPY

Principal

Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



# Nonlocal Initial Value Problems For First-Order Dynamic Equations On Time Scales\*

Sanket Tikare<sup>†</sup>

Received 5 June 2020

## Abstract

This article is concerned with a class of first-order dynamic equations on time scales with nonlocal initial conditions. Qualitative and quantitative results are discussed. Through an application of a fixed point theorem due to O'Regan, the existence of solutions is investigated. Under suitable assumptions, we deduce the existence result for nonlocal dynamic Cauchy problem. We also examine the continuous dependency of solutions on initial conditions. We illustrate our main result through examples.

## 1 Introduction

The theory of time scales calculus, introduced by S. Hilger [15], is an efficient tool to unify both continuous and discrete problems in one theory. The advantage of working on dynamic equations on time scales is that, under one framework, we can describe continuous-discrete hybrid processes. The results obtained, in the context of time scales, are more general and includes various other results as a particular case. Thus, the study of dynamic equations on time scales has potential applications to any field involving both continuous and discrete dynamical processes. As a consequence, in recent decades, the field of dynamic equations on time scales has become very popular to be used in mathematical modelling of several situations where simultaneous modelling is needed.

The first-order dynamic equations on time scales have been studied extensively, covering a variety of different problems; see for instance [8, 16, 21, 24] and the references therein. Moreover, there has been significant growth in the study of initial and boundary value dynamic problems, see for instance [9, 10, 11, 23], but none of them considers nonlocal conditions. The study of nonlocal initial value problems constitutes a very interesting and important class of problems; because, in many physical systems, the measurements by a nonlocal condition may be more precise than the measurement given by a local initial condition. In the literature, a great deal of attention has been given to nonlocal problems for differential equations rather than for difference or dynamic equations, see [6, 7, 13, 14] and references therein. Thus, it is worthwhile to study dynamic equations on time scales with nonlocal initial conditions.

In this paper, we are interested in establishing the existence of solutions to the first-order dynamic equation of the form

$$x^\Delta(t) + p(t)x^\sigma(t) = f(t, x(t)), \quad t \in [0, T]_{\mathbb{T}} = [0, T] \cap \mathbb{T}, \quad (1)$$

subject to the initial condition

$$x(0) = \Phi(x), \quad (2)$$

where  $T \in \mathbb{R}^+$ ,  $f : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous,  $\Phi : C([0, T]_{\mathbb{T}}, \mathbb{R}) \rightarrow \mathbb{R}$ , and  $p : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}$  is a function which is regressive and rd-continuous on  $\mathbb{T}$ . If  $\mathbb{T} = \mathbb{R}$ , then (1) becomes the well known ordinary differential equation [18]. If  $\mathbb{T} = \mathbb{Z}$ , then (1) becomes familiar difference equation [17].

Also, we discuss herein, the existence of solutions to the adjoint equation of (1)

$$x^\Delta(t) + q(t)x(t) = g(t, x(t)), \quad t \in [0, T]_{\mathbb{T}} = [0, T] \cap \mathbb{T}, \quad (3)$$

\*Mathematics Subject Classifications: 34N05; 34A12; 39A12.

<sup>†</sup>Department of Mathematics, Ramniranjan Jhunjhunwala College, Ghatkopar (W), Mumbai (M. S.) - 400 086, India



subject to (2), where  $q : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}$  is regressive and rd-continuous function,  $g : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous function.

Among various techniques available in the analysis that deals with the existence of solutions of dynamic problems, the method of fixed points is more elegant and powerful. Many authors have been employed several fixed point theorems in their investigation of the existence of solutions, see for instance [6, 7, 10, 13, 14, 22, 24]. For more details of fixed point theory, see [2, 12, 25].

In the present paper, we have applied a fixed point theorem of O'Regan, which is an extension of Krasnoselskii's fixed point theorem [25], to obtain the existence of solutions of dynamic problem (1)–(2).

The paper is organized as follows. In Section 2, we review some basic definitions and results needed in the paper. In Section 3, we present the results of the existence and continuous dependence of solutions to a nonlocal dynamic problem (1)–(2). Examples are provided in Section 4 to illustrate the main result. Finally, in Section 5 we make our concluding remarks.

## 2 Preliminaries

In this section, we recall some preliminary definitions and results which helps the reader to follow the paper easily. A nonempty closed subset of  $\mathbb{R}$ , which inherits the standard topology of  $\mathbb{R}$  is a time scale. It is denoted by  $\mathbb{T}$ . We assume that the reader of this paper is familiar with basic concepts of time scales calculus; these preliminary materials are very common in the literature and we refer the reader to [3, 4] for an excellent review of the topic. We begin by recalling briefly some necessary material from time scales calculus which are needed to follow the paper.

**Definition 1** A function  $x : \mathbb{T} \rightarrow \mathbb{R}$  is said to be rd-continuous if it is continuous at every right-dense points in  $\mathbb{T}$  and its left sided limits exist at left-dense points in  $\mathbb{T}$ .

**Definition 2** A function  $f : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$  is said to be rd-continuous on  $\mathbb{T} \times \mathbb{R}$  if  $f(\cdot, x)$  is rd-continuous on  $\mathbb{T}$  for each fixed  $x \in \mathbb{R}$  and  $f(t, \cdot)$  is continuous on  $\mathbb{R}$  for each fixed  $t \in \mathbb{T}$ .

**Definition 3** A function  $p : \mathbb{T} \rightarrow \mathbb{R}$  is said to be regressive if  $1 + \mu(t)p(t) \neq 0$  for all  $t \in \mathbb{T}^\kappa$ , where  $\mu : \mathbb{T} \rightarrow [0, \infty)$  is the graininess function defined by  $\mu(t) := \sigma(t) - t$ .

**Definition 4** For a regressive function  $p$ , the exponential function  $e_p(\cdot, t_0)$  on the time scale  $\mathbb{T}$  is defined as

$$e_p(t, t_0) := \exp \left( \int_{t_0}^t \xi_{\mu(s)}(p(s)) \Delta s \right) \text{ with } \xi_h(z) = \begin{cases} \frac{\text{Log}(1 + hz)}{h} & \text{if } h > 0, \\ z & \text{if } h = 0, \end{cases}$$

where  $\text{Log}$  is the principal logarithm function. In fact,  $e_p(\cdot, t_0)$  is the unique solution of the dynamic initial value problem

$$x^\Delta(t) = p(t)x, \quad x(t_0) = 1, \quad t, t_0 \in \mathbb{T},$$

where  $p$  is regressive and rd-continuous function.

For regressive functions  $p, q : \mathbb{T} \rightarrow \mathbb{R}$ , we define

$$p \oplus q := p + q + \mu pq, \quad \ominus p := \frac{-p}{1 + \mu p}, \quad p \ominus q := p \oplus (\ominus q).$$

Throughout the paper, we denote

$$E_0 := \sup_{t \in [0, T]_{\mathbb{T}}} |e_{\ominus p}(t, 0)| \text{ and } E := \sup_{s, t \in [0, T]_{\mathbb{T}}} |e_{\ominus p}(t, s)|.$$

Some fundamental properties of the exponential function are stated below.

**Certified as  
TRUE COPY**



**Principal**  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



**Theorem 1** Assume that  $p, q : \mathbb{T} \rightarrow \mathbb{R}$  are regressive and rd-continuous. Then the following hold.

- (i)  $e_0(t, s) \equiv 1$  and  $e_p(t, t) \equiv 1$ ;
- (ii)  $e_p(\sigma(t), s) = (1 + \mu(t)p(t))e_p(t, s)$ ;
- (iii)  $1/e_p(t, s) = e_{\ominus p}(t, s)$ ;
- (iv)  $e_p(t, s) = 1/e_p(s, t)$ ;
- (v)  $e_p(t, s)e_p(s, r) = e_p(t, r)$ ;
- (vi)  $e_p(t, s)e_q(t, s) = e_{p \oplus q}(t, s)$ ;
- (vii)  $e_p(t, s)/e_q(t, s) = e_{p \ominus q}(t, s)$ .

We now remind of essential features from the fixed point theory. Let  $C([0, T]_{\mathbb{T}}, \mathbb{R})$  be the family of all continuous functions from  $[0, T]_{\mathbb{T}}$  into  $\mathbb{R}$ , which is Banach space coupled with the norm  $\|\cdot\|$  defined as  $\|x\| := \sup_{t \in [0, T]_{\mathbb{T}}} |x(t)|$ . We denote this Banach space by  $X$  and a closed ball of radius  $r$  centered in the null element of  $X$  by  $B_r$ ; that is,  $B_r := \{x \in X : \|x\| \leq r\}$ .

**Definition 5 ([12])** Let  $X, Y$  be two Banach spaces. A mapping  $F : X \rightarrow Y$  is said to be completely continuous if the image of each bounded set  $B$  of  $X$ ,  $F(B)$ , is relatively compact in  $Y$ .

**Definition 6 ([19])** Let  $X$  be a Banach space. A mapping  $F : X \rightarrow X$  is said to be nonlinear contraction map if there exists a continuous, nondecreasing function  $\phi : [0, \infty) \rightarrow [0, \infty)$  satisfying  $\phi(z) < z$  for  $z > 0$  such that

$$\|F(x) - F(y)\| \leq \phi(\|x - y\|)$$

for all  $x, y \in X$ .

The version of Arzelà-Ascoli theorem in the context of time scales is stated as follows, see [1, Lemma 2.6].

**Theorem 2** A bounded equicontinuous subset  $D$  of  $C(\mathbb{T}, \mathbb{R})$  is relatively compact.

The main result of this paper is based upon an application of the following fixed point theorem of D. O'Regan [19].

**Theorem 3** Let  $U$  be an open set in a closed, convex set  $C$  of a Banach space  $X$ . Assume  $0 \in U$ ,  $F(\bar{U})$  bounded and  $F : \bar{U} \rightarrow C$  is given by  $F := F_1 + F_2$ , where  $F_1 : \bar{U} \rightarrow X$  is continuous and completely continuous and  $F_2 : \bar{U} \rightarrow X$  is a nonlinear contraction. Then either,

- (A1)  $F$  has a fixed point in  $\bar{U}$ ; or
- (A2) there is a point  $u \in \partial U$  and  $\lambda \in (0, 1)$  with  $u = \lambda F(u)$ .

### 3 Main Results

In this section, we establish a result concerning the existence of solutions to (1)–(2) using Theorem 3 and its applications.

First, we present an auxiliary lemma which reformulate our dynamic problem (1)–(2) as equivalent delta integral equation. The idea is same as that of Lemma 3.1 [5]. Hence the proof is omitted.

Certified as  
TRUE COPY

  
Principal

Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



**Lemma 1** Let  $p : \mathbb{T} \rightarrow \mathbb{R}$  be rd-continuous and regressive. Suppose  $f : \mathbb{T} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous,  $t_0 \in \mathbb{T}$ , and  $\Phi : C(\mathbb{T}, \mathbb{R}) \rightarrow \mathbb{R}$ . Then  $x$  is the unique solution of the nonlocal dynamic problem (1)–(2) on  $\mathbb{T}$  if and only if

$$x(t) = e_{\ominus p}(t, t_0)\Phi(x) + \int_{t_0}^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \quad (4)$$

for all  $t \in \mathbb{T}$ .

**Theorem 4** Assume that  $f : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous function such that for all  $t \in [0, T]_{\mathbb{T}}$  and for each  $x \in B_r$ ,  $|f(t, x)| \leq h_r(t)$ , where  $h_r : [0, T]_{\mathbb{T}} \rightarrow [0, \infty)$  is such that

$$\lim_{r \rightarrow \infty} \left( \frac{1}{r} \int_0^T h_r(t)\Delta t \right) = \beta < \frac{1}{E} < \infty. \quad (5)$$

Also, suppose that there is a continuous, nondecreasing function  $m : [0, \infty) \rightarrow [0, \infty)$  satisfying  $m(z) < z$  for  $z > 0$  such that

$$|\Phi(x) - \Phi(y)| \leq \frac{1}{E_0} m(\|x - y\|) \text{ for all } x, y \in X. \quad (6)$$

Then the dynamic nonlocal problem (1)–(2) has at least one solution.

**Proof.** Select  $r > 0$  such that for every  $x \in X$ ,

$$r > E_0|\Phi(x)| + E \int_0^T h_r(s)\Delta s.$$

Then  $B_r$  is a closed, convex subset of  $X$  and  $0 \in B_r$ . Here 0 is a null element of  $X$ . We define the mapping  $F : B_r \rightarrow X$  by

$$F(x)(t) := F_1(x)(t) + F_2(x)(t), \quad t \in [0, T]_{\mathbb{T}}, \quad (7)$$

where  $F_i : B_r \rightarrow X$  ( $i = 1, 2$ ) are given by

$$F_1(x)(t) := \int_0^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \quad (8)$$

and

$$F_2(x)(t) := e_{\ominus p}(t, 0)\Phi(x). \quad (9)$$

Obviously, the fixed points of  $F$  are the solutions of (1)–(2). We verify that the conditions of Theorem 3 are fulfilled.

1)  $F(B_r)$  is bounded. Let  $x \in B_r$  be arbitrary. Then for each  $t \in [0, T]_{\mathbb{T}}$ ,

$$F(x)(t) = e_{\ominus p}(t, 0)\Phi(x) + \int_0^t e_{\ominus p}(t, s)f(s, x(s))\Delta s, \quad (10)$$

and we see that

$$\begin{aligned} |F(x)(t)| &\leq \sup_{t \in [0, T]_{\mathbb{T}}} |F(x)(t)| \\ &= \sup_{t \in [0, T]_{\mathbb{T}}} \left| e_{\ominus p}(t, 0)\Phi(x) + \int_0^t e_{\ominus p}(t, s)f(s, x(s))\Delta s \right| \\ &\leq \sup_{t \in [0, T]_{\mathbb{T}}} \left\{ |e_{\ominus p}(t, 0)| |\Phi(x)| + \int_0^t |e_{\ominus p}(t, s)| |f(s, x(s))| \Delta s \right\} \\ &\leq E_0 |\Phi(x)| + E \int_0^T h_r(s)\Delta s \\ &< r. \end{aligned}$$

Certified as  
TRUE COPY



Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



Thus,  $\|F(x)\| < r$  and hence  $F(B_r)$  is bounded.

2)  $F_1 : B_r \rightarrow X$  is continuous and completely continuous. Let  $(x_n)$  be a sequence of elements of  $B_r$  converges to  $x$  in  $B_r$ . Then we see that,

$$\begin{aligned} \|F_1(x_n) - F_1(x)\| &= \sup_{t \in [0, T]_{\mathbb{T}}} |F_1(x_n)(t) - F_1(x)(t)| \\ &= \sup_{t \in [0, T]_{\mathbb{T}}} \left| \int_0^t e_{\ominus p}(t, s) f(s, x_n(s)) \Delta s - \int_0^t e_{\ominus p}(t, s) f(s, x(s)) \Delta s \right| \\ &\leq \sup_{t \in [0, T]_{\mathbb{T}}} \left\{ \int_0^t |e_{\ominus p}(t, s)| |f(s, x_n(s)) - f(s, x(s))| \Delta s \right\} \\ &\leq E \int_0^T |f(s, x_n(s)) - f(s, x(s))| \Delta s, \end{aligned}$$

which yields by rd-continuity of  $f$  that

$$\|F_1(x_n) - F_1(x)\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus,  $F_1$  is uniformly continuous on  $B_r$  and hence is continuous on  $B_r$ .

Next, we show that  $F_1$  is completely continuous. For this, we show that for each bounded subset  $D$  of  $B_r$ ,  $F_1(D)$  is in  $D$ . We claim that there exists a positive number  $r_1$  such that  $F_1(B_{r_1}) \subseteq B_{r_1}$ . Suppose not, then we see that for each positive number  $r$  there is  $x \in B_r$  satisfying  $|F_1(x)(t)| > r$  for all  $t \in [0, T]_{\mathbb{T}}$ . Now

$$\begin{aligned} r &< |F_1(x)(t)| \\ &= \left| \int_0^t e_{\ominus p}(t, s) f(s, x(s)) \Delta s \right| \\ &\leq \int_0^t |e_{\ominus p}(t, s)| |f(s, x(s))| \Delta s \\ &\leq \sup_{t \in [0, T]_{\mathbb{T}}} \left\{ \int_0^t |e_{\ominus p}(t, s)| |f(s, x(s))| \Delta s \right\} \\ &\leq E \int_0^T h_r(s) \Delta s. \end{aligned}$$

Thus,

$$1 < \frac{E}{r} \int_0^T h_r(s) \Delta s.$$

Taking limit as  $r \rightarrow \infty$  on both sides, we obtain  $1 < E\beta$ , which is contradiction. Hence there exists a positive number  $r_1$  such that  $F_1(B_{r_1}) \subseteq B_{r_1}$ . Let  $t_1, t_2 \in [0, T]_{\mathbb{T}}$  with  $t_1 < t_2$  and fix  $x \in B_r$ . Then we compute that

$$\begin{aligned} &|F_1(x)(t_2) - F_1(x)(t_1)| \\ &= \left| \int_0^{t_2} e_{\ominus p}(t_2, s) f(s, x(s)) \Delta s - \int_0^{t_1} e_{\ominus p}(t_1, s) f(s, x(s)) \Delta s \right| \\ &= \left| e_{\ominus p}(t_2, 0) \int_0^{t_1} e_p(s, 0) f(s, x(s)) \Delta s + e_{\ominus p}(t_2, 0) \int_{t_1}^{t_2} e_p(s, 0) f(s, x(s)) \Delta s \right. \\ &\quad \left. - e_{\ominus p}(t_1, 0) \int_0^{t_1} e_p(s, 0) f(s, x(s)) \Delta s \right| \\ &\leq |e_{\ominus p}(t_2, 0) - e_{\ominus p}(t_1, 0)| \int_0^{t_1} |e_p(s, 0)| |f(s, x(s))| \Delta s \\ &\quad + |e_{\ominus p}(t_2, 0)| \int_{t_1}^{t_2} |e_p(s, 0)| |f(s, x(s))| \Delta s. \end{aligned}$$

Certified as  
TRUE COPY



Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



Since  $e_{\Theta p}(t, \cdot)$  is continuous on  $[0, T]_{\mathbb{T}}$ , the right hand side tends to zero as  $t_1 \rightarrow t_2$  or  $t_2 \rightarrow t_1$ . Thus, by definition 5 and Theorem 2, we obtain that  $F_1$  is completely continuous. Hence  $F_1 : B_r \rightarrow X$  is continuous and completely continuous.

3)  $F_2 : B_r \rightarrow X$  is nonlinear contraction map. For  $x, y \in B_r$  and for each  $t \in [0, T]_{\mathbb{T}}$ , consider

$$F_2(x)(t) - F_2(y)(t) = e_{\Theta p}(t, 0)\Phi(x) - e_{\Theta p}(t, 0)\Phi(y).$$

Then we see that

$$\begin{aligned} \|F_2(x)(t) - F_2(y)(t)\| &= \sup_{t \in [0, T]_{\mathbb{T}}} |F_2(x)(t) - F_2(y)(t)| \\ &= \sup_{t \in [0, T]_{\mathbb{T}}} |e_{\Theta p}(t, 0)| |\Phi(x) - \Phi(y)| \\ &\leq E_0 |\Phi(x) - \Phi(y)| \\ &\leq m(\|x - y\|). \end{aligned}$$

Thus, we have

$$|F_2(x)(t) - F_2(y)(t)| \leq m(\|x - y\|),$$

where  $m : [0, \infty) \rightarrow [0, \infty)$  is a continuous nondecreasing function satisfying  $m(z) < z$  for  $z > 0$ . Hence, in view of definition 6,  $F_2$  is nonlinear contraction map.

4) There is no  $u \in \partial B_r$  such that  $u = \lambda F(u)$  for  $\lambda \in (0, 1)$ . Suppose there is such a  $u \in \partial B_r$ . Then for each  $t \in [0, T]_{\mathbb{T}}$ ,

$$u(t) = \lambda F_1(u)(t) + \lambda F_2(u)(t).$$

We compute that,

$$\begin{aligned} \|u\| &= \sup_{t \in [0, T]_{\mathbb{T}}} |u(t)| \\ &\leq \sup_{t \in [0, T]_{\mathbb{T}}} \left| e_{\Theta p}(t, 0)\Phi(u) + \int_0^t e_{\Theta p}(t, s)f(s, u(s))\Delta s \right| \\ &\leq \sup_{t \in [0, T]_{\mathbb{T}}} \left\{ |e_{\Theta p}(t, 0)| |\Phi(u)| + \int_0^t |e_{\Theta p}(t, s)| |f(s, u(s))| \Delta s \right\} \\ &\leq E_0 |\Phi(u)| + E \int_0^T h_r(s)\Delta s. \end{aligned}$$

Since  $u \in \partial B_r$ , we obtain

$$r \leq E_0 |\Phi(u)| + E \int_0^T h_r(s)\Delta s,$$

which contradicts to the choice of  $r > 0$ . Hence there is no  $u \in \partial B_r$  for which  $u = \lambda F(u)$  for  $\lambda \in (0, 1)$ . Consequently, we can deduce from Theorem 3 that  $F$  has a fixed point in  $B_r$ . Hence there exists at least one solution of (1)–(2). This completes the proof. ■

Now, we shall turn our attention to the existence of solutions to the dynamic nonlocal problem (3)–(2). Notice that the simple useful formula transform (3) into (1). Hence, we will use Theorem 4 to formulate the existence result for dynamic nonlocal problem (3)–(2).

**Theorem 5** Consider the dynamic nonlocal problem (3)–(2). Let  $q$  be rd-continuous, regressive function such that  $1 - \mu(t)q(t) > 0$  for all  $t \in [0, T]_{\mathbb{T}}$ . Assume that  $g : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous function such that for all  $t \in [0, T]_{\mathbb{T}}$  and for each  $x \in B_r$ ,

$$|g(t, x)| \leq h_r(t)(1 - \mu q(t)), \quad (11)$$

Certified as  
TRUE COPY



Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



where the functions  $h_r$  and  $\Phi$  are defined as in Theorem 4. Then the dynamic nonlocal problem (3)–(2) has at least one solution.

**Proof.** Replacing  $x$  in (3) by the simple useful formula  $x^\sigma := x + \mu x^\Delta$ , we obtain

$$x^\Delta(t) + \left( \frac{q(t)}{1 - \mu q(t)} \right) x^\sigma(t) = \frac{g(t, x(t))}{1 - \mu q(t)}.$$

This equation is essentially in the form (1) with  $p(t) := \frac{q(t)}{1 - \mu q(t)}$  and  $f(t, x) := \frac{g(t, x)}{1 - \mu q(t)}$ . Also, we notice that  $p$  is regressive and rd-continuous and  $f$  is rd-continuous. Now, it is not difficult to show that (11) reduces to  $|f(t, x)| \leq h_r(t)$  for all  $t \in [0, T]_{\mathbb{T}}$  and for each  $x \in B_r$ . Hence the result follows from Theorem 4. ■

Next, again Theorem 4 will be used to guarantee the existence of at least one solution to the nonlocal dynamic Cauchy problem of the type

$$x^\Delta(t) = F(t, x), \quad t \in [0, T]_{\mathbb{T}} \quad (12)$$

subject to (2), where  $F : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous and possibly nonlinear function.

**Theorem 6** Consider the dynamic nonlocal Cauchy problem (12)–(2). Let  $q$  be rd-continuous, regressive function such that  $1 - \mu(t)q(t) > 0$  for all  $t \in [0, T]_{\mathbb{T}}$ . Assume that  $F : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  is rd-continuous function such that for all  $t \in [0, T]_{\mathbb{T}}$  and for each  $x \in B_r$ ,

$$|F(t, x) + q(t)x| \leq h_r(t)(1 - q(t)x),$$

where the functions  $h_r$  and  $\Phi$  are defined as in Theorem 4. Then the dynamic nonlocal Cauchy problem (12)–(2) has at least one solution.

**Proof.** We write (12) as

$$x^\Delta(t) = -q(t)x^\sigma(t) + q(t)x^\sigma(t) + F(t, x(t)),$$

which by using so-called simple useful formula yields that

$$x^\Delta(t) = \frac{-q(t)}{1 - \mu(t)q(t)} x^\sigma(t) + \frac{q(t)x(t) + F(t, x(t))}{1 - \mu(t)q(t)}.$$

Let  $p : [0, T]_{\mathbb{T}} \rightarrow \mathbb{R}$  be defined by

$$p(t) := \frac{q(t)}{1 - \mu(t)q(t)}.$$

Then clearly,  $p$  is rd-continuous and regressive. Again, define  $f : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$f(t, x) := \frac{q(t)x + F(t, x)}{1 - \mu(t)q(t)}$$

which is rd-continuous. With these notations, (12) may be written as

$$x^\Delta(t) + p(t)x^\sigma(t) = f(t, x(t)) \quad (13)$$

which is essentially (1). Now for all  $t \in [0, T]_{\mathbb{T}}$  and for each  $x \in B_r$ ,

$$|f(t, x)| = \frac{|q(t)x + F(t, x)|}{1 - \mu(t)q(t)} \leq \frac{h_r(t)(1 - \mu(t)q(t))}{1 - \mu(t)q(t)}.$$

Hence  $|f(t, x)| \leq h_r(t)$ . Thus, we are able to apply Theorem 4 to the dynamic nonlocal problem (13)–(2) and obtain that the dynamic nonlocal Cauchy problem (12)–(2) has at least one solution. This completes the proof. ■

Certified as  
TRUE COPY



Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



**Remark 1** We remark that Theorems 4, 5, and 6 also hold and can be proved by using the same analysis, even if  $f$  is  $\Delta$ -Carathéodory function. In this case, we get  $\Delta$ -Carathéodory type solutions. For such solutions, the dynamic nonlocal problem (1)–(2) is equivalent to the integral equation (4)  $\Delta$ - a.e.  $t \in [0, T]_{\mathbb{T}}$ . For more details of  $\Delta$ -Carathéodory function and solutions, the readers are referred to [11, 20].

The dependency of solutions to (1)–(2) with respect to the initial conditions can be obtained as follows.

**Theorem 7** Suppose  $f : [0, T]_{\mathbb{T}} \times \mathbb{R} \rightarrow \mathbb{R}$  and  $\Phi_1, \Phi_2 : X \rightarrow \mathbb{R}$  satisfy the hypotheses of Theorem 4. Also, suppose for all  $t \in [0, T]_{\mathbb{T}}$  and for all  $x, y \in X$ ,

$$|f(t, x) - f(t, y)| \leq \frac{1}{E} \|x - y\|. \tag{14}$$

Then for the corresponding solutions  $x_1, x_2$  of the dynamic equation (1) subject to the initial conditions

$$x(0) = \Phi_i(x) \quad (i = 1, 2), \tag{15}$$

the inequality

$$\|x_1 - x_2\| \leq E_0 |\Phi_1(x_1) - \Phi_2(x_2)| \sup_{t \in [0, T]_{\mathbb{T}}} |e_1(t, 0)|.$$

holds. Additionally, if  $|\Phi_1(x_1) - \Phi_2(x_2)| \leq \delta$  for some  $\delta > 0$ , then we have

$$\|x_1 - x_2\| \leq E_0 \delta \sup_{t \in [0, T]_{\mathbb{T}}} |e_1(t, 0)|. \tag{16}$$

**Proof.** Let  $x_i$  ( $i = 1, 2$ ) be the corresponding solutions to the dynamic problem (1)–(15). Then from Lemma 1, these solutions are given by

$$x_i(t) = e_{\Theta p}(t, 0)\Phi_i(x_i) + \int_0^t e_{\Theta p}(t, s)f(s, x_i(s))\Delta s \quad (i = 1, 2). \tag{17}$$

Therefore, for each  $t \in [0, T]_{\mathbb{T}}$ , we obtain

$$\begin{aligned} |x_1(t) - x_2(t)| &= \left| e_{\Theta p}(t, 0)\Phi_1(x_1) + \int_0^t e_{\Theta p}(t, s)f(s, x_1(s))\Delta s \right. \\ &\quad \left. - e_{\Theta p}(t, 0)\Phi_2(x_2) + \int_0^t e_{\Theta p}(t, s)f(s, x_2(s))\Delta s \right| \\ &\leq |e_{\Theta p}(t, 0)| |\Phi_1(x_1) - \Phi_2(x_2)| + \int_0^t |e_{\Theta p}(t, s)| |f(s, x_1(s)) - f(s, x_2(s))| \Delta s \\ &\leq E_0 |\Phi_1(x_1) - \Phi_2(x_2)| + E \int_0^t |f(s, x_1(s)) - f(s, x_2(s))| \Delta s. \end{aligned}$$

By inequality (14), we can write

$$|x_1(t) - x_2(t)| \leq E_0 |\Phi_1(x_1) - \Phi_2(x_2)| + \int_0^t |x_1(s) - x_2(s)| \Delta s. \tag{18}$$

Now for  $t \in [0, T]_{\mathbb{T}}$ , let

$$F(t) = \int_0^t |x_1(s) - x_2(s)| \Delta s.$$

Then, we see from (18) that

$$F^\Delta(t) - F(t) \leq E_0 |\Phi_1(x_1) - \Phi_2(x_2)|.$$

**Certified as  
TRUE COPY**



**Principal**  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



Dividing both sides by  $e_1(\sigma(t), 0)$  and integrating from 0 to  $t$ ,  $t \in [0, T]_{\mathbb{T}}$ , we obtain

$$e_1(0, t)F(t) \leq E_0|\Phi_1(x_1) - \Phi_2(x_2)|[e_1(0, 0) - e_1(0, t)],$$

which yields that

$$F(t) \leq E_0|\Phi_1(x_1) - \Phi_2(x_2)|[e_1(t, 0) - 1].$$

Now (18) becomes

$$\begin{aligned} |x_1(t) - x_2(t)| &\leq E_0|\Phi_1(x_1) - \Phi_2(x_2)| + E_0|\Phi_1(x_1) - \Phi_2(x_2)|[e_1(t, 0) - 1] \\ &= E_0|\Phi_1(x_1) - \Phi_2(x_2)|e_1(t, 0). \end{aligned}$$

Thus,

$$\|x_1 - x_2\| \leq E_0|\Phi_1(x_1) - \Phi_2(x_2)| \sup_{t \in [0, T]_{\mathbb{T}}} |e_1(t, 0)|.$$

Finally, inequality (16) is consequence of the above inequality. This completes the proof. ■

## 4 Examples

In this section, we present two examples to complement and expound our main result. Throughout this section, we take  $\mathbb{T} = [0, 1] \cup [2, 3]$ .

Consider the dynamic problem (1)–(2) on  $[0, 3]_{\mathbb{T}}$ .

(a) In (1)–(2), we take  $p \equiv 1$ ,  $f(t, x) = \frac{x^3}{x^2 + 3} + t$ , and  $\Phi(x) = \sqrt{1 + x}$ .

Then for  $x \in B_r$ , we see that  $|f(t, x)| \leq r + t$ . Also, we get  $\beta = 1$ .

Now, for  $x, y \in B_r$ ,

$$|\Phi(x) - \Phi(y)| \leq 1 + \sqrt{\|x - y\|}.$$

We take  $m(z) = 1 + \sqrt{z}$ . Then  $m$  is decreasing function such that

$$|\Phi(x) - \Phi(y)| \leq m(\|x - y\|).$$

It is easy to see that  $E = 1/e$  and  $E\beta < 1$ . From Theorem 4, we conclude that the dynamic problem (1)–(2) has a solution on  $[0, 3]_{\mathbb{T}}$ .

(b) We can take  $p \equiv 1$ ,  $f(t, x) = \frac{x^2}{x^2 + 3} + t$ , and  $\Phi(x) = x^{2/3}$ . Then we see that  $\beta = 0$  and the conditions of Theorem 4 are satisfied. Thus the dynamic problem (1)–(2) has a solution on  $[0, 3]_{\mathbb{T}}$ .

## 5 Conclusions

Throughout this paper, we extended the concept of nonlocal initial value problems to time scales. We obtained certain results concerning the existence of solutions; and discussed the continuous dependence of solutions to the initial conditions. Moreover, we complemented and expounded our result through examples. We believe that, in the context of general time scales, the results presented here are promising in simultaneous modelling of both continuous and discrete processes. Further, it is possible to extend the results of this paper easily to systems of equations and Banach spaces. Also, as a continuation of this work, the other aspects like the approximation of solutions, boundedness of solutions, upper and lower solutions can be studied in the near future.

**Acknowledgments.** The author expresses his gratitude to the referees for their insightful comments and suggestions which helped to improve the quality of the paper. The author is also thankful to Prof. Christopher C. Tisdell for his valuable suggestions towards the presentation of the paper. This work was supported by the Seed Grant of Ramniranjan Jhunjunwala College (Ref. No. 935-17102019).

Certified as  
TRUE COPY




Principal  
Ramniranjan Jhunjunwala College,  
Ghatkopar (W), Mumbai-400086.



## References

- [1] R. P. Agarwal, M. Bohner, and P. Řehák, Half-linear Dynamic Equations. Nonlinear Analysis and Applications: To V. Lakshmikantham on his 80th birthday. Vol. 1, 2, 1–57, Kluwer Acad. Publ., Dordrecht, 2003.
- [2] R. P. Agarwal, M. Meehan, and D. O'Regan, Fixed Point Theory and Applications, Cambridge University Press, Cambridge, 2004.
- [3] M. Bohner and A. C. Peterson, Dynamic Equations on Time Scales: An Introduction with Applications, Birkhäuser, Boston, 2001.
- [4] M. Bohner and A. C. Peterson, Advances in Dynamic Equations on Time Scales, Birkhäuser, Boston, 2003.
- [5] M. Bohner, S. Tikare and I. L. D. Santos, First-order nonlinear dynamic initial value problems, Int. J. Dyn. Syst. Differ. Equ., 2021, To appear.
- [6] A. Boucherif, Differential equations with nonlocal boundary conditions, Nonlinear Anal., 47(2001), 2419–2430.
- [7] A. Boucherif and R. Precup, On the nonlocal initial value problem for first order differential equations, Fixed Point Theory, 4(2003), 205–212.
- [8] L. Bourlin and E. Trélat, General Cauchy–Lipschitz theory for  $\Delta$ -cauchy problems with Carathéodory dynamics on time scales, J. Differ. Equ. Appl., 20(2014), 526–547.
- [9] A. Cabada and D. R. Vivero, Existence of solutions of first-order dynamic equations with nonlinear functional boundary value conditions, Nonlinear Anal., 63(2005), e697–e706.
- [10] Q. Dai and C. C. Tisdell, Existence of solutions to first-order dynamic boundary value problems, Int. J. Difference Equ., 1(2006), 1–17.
- [11] H. Gilbert, Existence Theorems for first-order equations on time scales with  $\Delta$ -Carathéodory functions, Adv. Difference Equ., 2010, Art. ID 650827, 20 pp.
- [12] A. Granas and J. Dugundji, Fixed Point Theory, Springer–Verlag, New York, 2003.
- [13] E. M. A. Hamd–Allah, On the existence of solutions of two differential equations with a nonlocal condition, J. Egyptian Math. Soc., 24(2016), 367–372.
- [14] J. Henderson, Existence of local solutions for boundary value problems with integral conditions, Commun. Appl. Anal., 19(2015), 103–112.
- [15] S. Hilger, Analysis on measure chains—A unified approach to continuous and discrete calculus, Results Math., 18(1990), 18–56.
- [16] B. Karpuz, On the existence and uniqueness of solutions to dynamic equations, Turk. J. Math., 42(2018), 1072–1089.
- [17] W. G. Kelley and A. C. Peterson, Difference Equations, An Introduction with Applications, Second Edition, Academic Press, New York, 2001.
- [18] W. G. Kelley and A. C. Peterson, Differential Equations, Classical and Qualitative, Second Edition, Springer Science, New York, 2010.
- [19] D. O'Regan, Fixed-point theory for the sum of two operators, Appl. Math. Lett., 9(1996), 1–8.

**Certified as  
TRUE COPY**

  
**Principal**  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.



- [20] I. L. D. Santos, Discontinuous dynamic equations on time scales, *Rend. Circ. Mat. Palermo*, 64(2015), 383–402.
- [21] S. Tikare, Generalized first order dynamic equations on time scales with  $\Delta$ -Carathéodory functions, *Differ. Equ. Appl.*, 11(2019), 167–182.
- [22] S. Tikare and I. L. D. Santos, Existence results for solutions to discontinuous dynamic equations on time scales, *Differ. Equ. Appl.*, 12(2020), 89–100.
- [23] C. C. Tisdell, Existence and uniqueness to nonlinear dynamic boundary value problems, *Cubo*, 8(2006), 11–23.
- [24] C. C. Tisdell and A. Zaidi, Basic qualitative and quantitative results for solutions to nonlinear, dynamic equations on time scales with an application to economic modelling, *Nonlinear Anal.*, 68(2008), 3504–3524.
- [25] E. Zeidler, *Nonlinear Functional Analysis and its Applications. Fixed-Point Theorems*, Translated from the German by Peter R. Wadsack, Springer-Verlag, New York, 1986.

**Certified as  
TRUE COPY**



**Principal  
Ramniranjan Jhunjhunwala College,  
Ghatkopar (W), Mumbai-400086.**